

The Relationship between Maternal Haemoglobin and Haematocrit with Low Birth Weight and Preterm Labour

Hamid Kermani Moakhar (MD)¹, Behnaz Sovizi (MD)², Mitra Eftekhari Yazdi (MD)^{2*}

¹ Medical doctor, Sabzevar University of Medical Sciences, Sabzevar, Iran

² Assistant professor, Department of Obstetrics and Gynaecology, School of Medicine, Sabzevar University of Medical Sciences, Sabzevar, Iran

ARTICLE INFO

Article type:
Original article

Article History:
Received: 04- May-2018
Accepted: 29- Aug-2018

Key words:
Haemoglobin
Haematocrit
Low birth weight
Preterm labour
Anemia

ABSTRACT

Background & aim: Low levels of maternal haemoglobin and haematocrit in the first, second, and third trimesters are considered as possible risk factors regarding low birth weight (LBW) and preterm labour (PTL). The present study aimed to evaluate the relationship between maternal haemoglobin (Hb) and haematocrit (HCT) levels with LBW and PTL.

Methods: This cross-sectional study was conducted on 383 pregnant women who were admitted in postnatal ward and gave birth to live neonates. Maternal Hb and HCT levels in the first, second, and third trimesters were obtained from medical records. The data were analysed using chi-square test by SPSS software (version 19).

Results: A total of 383 pregnant women with the mean age of 25.5 years were participated in this study. There was a significant relationship between maternal HCT in the first, second, and third trimesters with LBW ($P < 0.01$). Also a significant relationship was observed between maternal Hb in the first, second, and third trimesters with the LBW ($P < 0.01$). In addition, a significant relationship was reported between maternal HCT with the PTL ($P < 0.01$). There was a significant correlation between maternal Hb in the first, second, and third trimesters with the PTL ($P = 0.01$).

Conclusion: Based on the findings of this study, it was suggested that maternal anaemia should be diagnosed and treated at any stage of pregnancy in order to reduce the risk of the LBW and PTL.

► Please cite this paper as:

Kermani Moakhar H, Sovizi B, Eftekhari Yazdi M. The Relationship between Maternal Haemoglobin and Haematocrit with Low Birth Weight and Preterm Labour. Journal of Midwifery and Reproductive Health. 2019; 7(1): 1577-1583. DOI: 10.22038/jmrh.2018.31621.1341

Introduction

Neonatal mortality rate is among the most important health indicators in every country, which is linked to the adequacy of maternal health care and social and economic factors (1). One of the determinants of neonatal mortality is low birth weight (LBW) (1, 2). The LBW is defined as birth weight of a newborn between 1500 and 2500 g (1). Mortality rate in the neonates with LBW is reported as 40 times and in newborns with very low birth weight (VLBW), between 1000 and 1500 g, is announced as 200 times higher than that in neonates with normal birth weight (2). According to statistics, 90% of the newborns

with LBW are annually born in developing countries (3). In some areas of Asia, one out of two live births is considered to be a neonate with LBW (3).

In Iran, 289 children below 5 years old daily die among which 48% are neonates mostly with LBW (3). Maternal age, occupation, and weight, as well as a history of LBW in previous pregnancies, the number of pregnancies along with addition and pregnancy duration have been identified as predicting factors for the LBW (4-6). On the other hand, poor socio-economic condition, malnutrition, hypertension, multiple pregnancies, and low maternal haemoglobin

* Corresponding author: Mitra Eftekhari Yazdi, Assistant professor, Department of Obstetrics and Gynaecology, School of Medicine, Sabzevar University of Medical Sciences, Sabzevar, Iran. Tel: +989155710054; Email: m_eftekhari1144@yahoo.com

(Hb) and haematocrit (HCT) levels, as well as illiteracy, maternal chronic diseases, and alcohol use were reported to be related with the LBW (7-9). It is believed that the LBW is associated with mental retardation, learning disorders, and cognitive growth retardation, as well as visual-motor disorders, and cerebral palsy (CP) (10).

Preterm labour (PTL) is defined as giving birth earlier than 37 weeks of gestation (11). Based on the differences observed in mortality and morbidity, the PTL was divided into the following categories based on the gestational age (GA), which were mild preterm (between 32 and 36 weeks of gestation), very preterm (between 28 and 31 weeks of gestation), and extremely preterm (earlier than 28 weeks of gestation) (12). The PTL may be related to cardiopulmonary disorders, brain growth retardation, and various anomalies (13). Morbidity and mortalities due to the PTL are mainly attributed to prematurity of the organs and include respiratory distress syndrome, bronchopulmonary dysplasia, pneumothorax, pneumonia, patent ductus arteriosus, intraventricular haemorrhage, periventricular leukomalacia, and the CP (13). The neonates with both LBW and PTL are at extremely high risks regarding morbidity and mortality (2, 13, 14).

Anaemia in pregnancy is defined as the HCT level below 33% or the Hb level below 11g/dl in the first and third trimester and the Hb below 10.5 g/dl in the second trimester (15, 16). Anaemia in pregnancy can lead to the PTL, intrauterine growth retardation, reduced physical activity capacity, immune response disorders, and heart failure in mother; however, it can result in the LBW, PTL, low Apgar score, and perinatal mortality in newborns (17, 18). It was previously reported that anaemia in the first half of pregnancy is related to increased risk of the PTL (19). Therefore, maternal anaemia is among the preventable causes of the PTL and LBW. On the other hand, the higher values of the HCT are related to maternal and neonatal adverse outcomes (19). The present study was carried out in order to evaluate the maternal factors, which contribute to the neonatal LBW and PTL in women that gave birth to live neonates and were admitted in the obstetric ward of Mobini Hospital, Sabzevar,

Iran.

Materials and Methods

This cross-sectional study was conducted on women that gave birth to live newborns in Mobini Hospital, Sabzevar, Iran in 2016. The inclusion criteria were giving birth to live infant and having complete medical records. The exclusion criteria were multiple pregnancies, history of diabetes, thalassemia or sickle cell anaemia in mother, and infants with gross congenital abnormalities, as well as pre-eclampsia and the obstetric causes of the PTL, including placenta abruption or premature rupture of membranes.

All the participants were interviewed by the author and asked to provide information regarding their age, infant gender, place of residence, educational level, the number of pregnancies, number of health care visits during pregnancy, and neonatal outcomes. Birth weight was obtained from medical records of the newborns and GA was obtained from the medical records based on ultrasounds. The Hb and HCT levels in the first, second, and third trimester were obtained from medical records of the subjects. The HB and HCT were measured by Technicon H-1 blood counter after taking peripheral venous blood in Ethylenediaminetetraacetic acid tubes. The HCT was divided into three groups, which were low (<34%), normal (34%-40%), and high (>40%), while the Hb was grouped into low (<10.5g/dl), normal (10.5-13g/dl), and high (>13 g/dl) based on the WHO criteria.

Statistical analysis

The data were analysed using SPSS (version 19). The Continuous data were checked for normality by Kolmogorov-Smirnov test. The mean and standard deviation were utilized to describe quantitative variables. The Qualitative variables were presented as frequency and percentage. The Chi-square test was used to evaluate the relation between categorical variables. P-values lower than 0.05 were statistically considered significant.

Results

A total of 383 pregnant women were recruited in this study. The mean age of the

subjects was 25.5 years. The body mass index mean of the subjects was 22.6 kg/m². Table 1 tabulates the demographic characteristics of the women. The mean birth weight of the newborns was 3180 g (birth weight range: 2100-4700g). The LBW was observed in 21 (5.5%) of the neonates, while the birth weight was reported between 2500-4000 g in 353 (92.2%) newborns; furthermore, 9 (2.3%) cases had the birth weight of more than 4000 g.

The mean GA of the subjects was 38 weeks (minimum GA: 33 weeks, maximum GA: 42 weeks). The PTL, term, and post-term pregnancies were observed in 38 (9.9%), 296 (77.3%), and 49 (12.8%) subjects, respectively.

The HCT of the subjects in the first trimester was reported as 38% (minimum HCT: 30%, maximum HCT: 47%), while the HCT in the second and third trimesters were 35.5% (minimum HCT: 26%, maximum HCT: 45.5%) and 34.5% (minimum HCT: 28%, maximum HCT: 41.8%). The HCT categories of the subjects are shown in Table 2. The Mean Hb of the subjects in the first trimester was 12.8 g/dl (minimum Hb: 9 g/dl, maximum Hb: 16 g/dl), while the mean Hb in the second and third trimesters were 12.07 g/dl (minimum Hb: 9.5 g/dl, maximum Hb: 14.2 g/dl) and 11.5 g/dl (minimum Hb: 8.6 g/dl, maximum Hb: 14 g/dl). Table 2 presents the HCT categories of the cases.

Table 1. Demographic characteristics of subjects

| Variable | | Frequency | Percentage |
|--------------------------|----------------------|-----------|------------|
| Number of pregnancies | 1 | 180 | 47.0% |
| | 2 | 94 | 24.5% |
| | 3 | 98 | 25.6% |
| | ≥4 | 11 | 2.9% |
| Place of residence | Urban | 277 | 72.3% |
| | Rural | 106 | 27.7% |
| Level of education | Primary school | 49 | 12.8% |
| | Junior high school | 181 | 47.3% |
| | High school | 133 | 34.7% |
| | Graduate | 20 | 5.2% |
| Economic status | Good | 55 | 14.4% |
| | Moderate | 303 | 79.1% |
| | Poor | 25 | 6.5% |
| Height category | ≤150cm | 99 | 25.8% |
| | >150cm | 284 | 74.0% |
| Body Mass Index category | ≤20kg/m ² | 41 | 10.7% |
| | >21kg/m ² | 342 | 89.3% |
| Newborn gender | Male | 193 | 50.4% |
| | Female | 190 | 49.6% |

cm: centimetre; kg: kilogram; m: meter

Table 2. Descriptive statistics for HCT categories in each trimester among subjects

| Variable | | First-trimester Frequency (%) | Second-trimester Frequency (%) | Third-trimester Frequency (%) |
|-----------|-------------|----------------------------------|-----------------------------------|----------------------------------|
| HCT (%) | <33 | 24 (6.3%) | 75 (19.6%) | 138 (36%) |
| | 33-40 | 270 (70.5%) | 297 (77.5%) | 239 (62.4%) |
| | >40 | 89 (23.2%) | 11 (2.9%) | 6 (1.6%) |
| Hb (g/dl) | <10.5g/dl | 12 (3.1%) | 23 (6.0%) | 66 (17.2%) |
| | 10.5-13g/dl | 195 (50.9%) | 337 (88.0%) | 302 (78.9%) |
| | >13g/dl | 176 (46%) | 23 (6.0%) | 15 (3.9%) |

HCT: haematocrit; Hb: Haemoglobin; g: gram; dl: decilitre

Table 3. Relation between birth weight categories and HCT and Hb categories in each trimester

| | Birth weight | | |
|--|--------------|------------|--------|
| | <2500g | 2500-4000g | >4000g |

| | | | | | |
|---------------------------|-----|-------------|------------------------|------------|---------|
| 1 st trimester | HCT | <33% | 3(12.5%) ^a | 21(87.5%) | 0(0.0%) |
| | | 33%-40% | 16(5.9%) | 251(93.0%) | 3(1.1%) |
| | | >40% | 2(2.2%) | 81(91.0%) | 6(6.7%) |
| | Hb | <10.5g/dl | 3(25.0%) ^d | 9(75.0%) | 0(0.0%) |
| | | 10.5-13g/dl | 13(6.7%) | 179(91.8%) | 3(1.5%) |
| | | >13g/dl | 5(2.8%) | 165(93.8%) | 6(3.4%) |
| 2 nd trimester | HCT | <33% | 10(13.3%) ^b | 65(86.7%) | 0(0.0%) |
| | | 33%-40% | 11(3.7%) | 277(93.3%) | 9(3.0%) |
| | | >40% | 0(0.0%) | 11(100.0%) | 0(0.0%) |
| | Hb | <10.5g/dl | 5(21.7%) ^e | 18(78.3%) | 0(0.0%) |
| | | 10.5-13g/dl | 16(4.7%) | 312(92.6%) | 9(2.7%) |
| | | >13g/dl | 0(0.0%) | 23(100.0%) | 0(0.0%) |
| 3 rd trimester | HCT | <33% | 3(12.5%) ^c | 21(87.5%) | 0(0.0%) |
| | | 33%-40% | 16(5.9%) | 251(93.0%) | 3(1.1%) |
| | | >40% | 2(2.2%) | 81(91.0%) | 6(6.7%) |
| | Hb | <10.5g/dl | 14(21.2%) ^f | 52(100.0%) | 0(0.0%) |
| | | 10.5-13g/dl | 7(2.3%) | 286(94.7%) | 9(3.0%) |
| | | >13g/dl | 0(0.0%) | 15(100.0%) | 0(0.0%) |

Table 4. Relation between gestational age at birth categories and HCT and Hb categories in each trimester

| Time | Variable | Gestational age at birth | | | |
|---------------------------|----------|--------------------------|------------------------|------------|-----------|
| | | <37 weeks | 37-40 weeks | >40 weeks | |
| 1 st trimester | HCT | <33% | 6(25.0%) ^a | 12(50.0%) | 6(25.0%) |
| | | 33%-40% | 24(8.9%) | 218(80.7%) | 28(10.4%) |
| | | >40% | 8(9.0%) | 66(74.2%) | 15(16.9%) |
| | Hb | <10.5g/dl | 3(25.0%) ^d | 9(75%) | 0(0.0%) |
| | | 10.5-13g/dl | 27(13.8%) | 146(74.9%) | 22(11.3%) |
| | | >13g/dl | 8(4.5%) | 141(80.1%) | 27(15.3%) |
| 2 nd trimester | HCT | <33% | 20(26.7%) ^b | 41(54.7%) | 14(18.6%) |
| | | 33%-40% | 16(5.4%) | 246(82.8%) | 35(11.8%) |
| | | >40% | 2(18.2%) | 9(81.8%) | 0(0.0%) |
| | Hb | <10.5g/dl | 8(34.8%) ^e | 12(52.2%) | 3(13.0%) |
| | | 10.5-13g/dl | 28(8.3%) | 263(78.0%) | 46(13.6%) |
| | | >13g/dl | 2(8.7%) | 21(91.3%) | 0(0.0%) |
| 3 rd trimester | HCT | <33% | 29(21.0%) ^c | 84(60.9%) | 25(18.1%) |
| | | 33%-40% | 9(3.8%) | 206(86.2%) | 24(10.0%) |
| | | >40% | 0(0.0%) | 6(100.0%) | 0(0.0%) |
| | Hb | <10.5g/dl | 22(33.3%) | 33(50.0%) | 11(16.7%) |
| | | 10.5-13g/dl | 16(5.3%) | 248(82.1%) | 38(12.6%) |
| | | >13g/dl | 0(0.0%) | 15(100.0%) | 0(0.0%) |

The results of the chi-square test revealed a significant relation between birth weight with the HCT and Hb levels in the first, second, and third trimesters ($P<0.001$). The association between the GA at birth with the HCT and Hb levels in pregnancy were significant based on the results of chi-square test ($P<0.01$) (Table 4). The relationship between birth weight and gestational age, along with other study parameters, including maternal age, maternal height, the number of pregnancies, maternal

education level, and neonate gender, as well as place of residence, and economic status are shown in Table 5. There was a significant relationship between the number of pregnancies and maternal educational level ($P<0.05$) based

Table 5. Relation between other study parameters, birth weight, and gestational age categories

| Variable | | Gestational age at birth | | | P-value | Birth weight | | | P-value |
|-----------------------|--------------------|--------------------------|-------------|-----------|---------|--------------|------------|---------|---------|
| | | <37 weeks | 37-40 weeks | >40 weeks | | <2500g | 2500-4000g | >4000g | |
| Maternal age | <20 years | 6(7.9%) | 56(73.3%) | 14(18.4%) | 0.28 | 3(3.9%) | 70(92.1%) | 3(3.9%) | 0.08 |
| | 20-34 years | 24(9.6%) | 196(78.1%) | 31(12.4%) | | 11(4.4%) | 234(93.2%) | 6(2.4%) | |
| | >35 years | 8(14.3%) | 44(78.6%) | 4(7.1%) | | 7(12.5%) | 49(87.5%) | 0(0.0%) | |
| Maternal height | <150 cm | 11(11.1%) | 72(72.7%) | 16(16.2%) | 0.42 | 4(4.0%) | 93(93.9%) | 2(2.0%) | 0.73 |
| | <151 cm | 27(9.5%) | 224(78.9%) | 33(11.6%) | | 17(6.0%) | 260(91.5%) | 7(2.5*) | |
| Number of pregnancies | 1 st | 15(8.3%) | 140(77.8%) | 25(13.9%) | 0.02* | 6(3.3%) | 169(93.9%) | 5(2.8%) | 0.04* |
| | 2 nd | 8(8.5%) | 71(75.5%) | 15(16.0%) | | 6(6.4%) | 87(92.6%) | 1(1.1%) | |
| | 3 rd | 12(12.2%) | 77(78.6%) | 9(9.2%) | | 6(6.1%) | 89(90.8%) | 3(3.1%) | |
| | ≥4 th | 3(27.3%) | 8(72.7%) | 0(0.0%) | | 3(27.3%) | 8(72.7%) | 0(0.0%) | |
| Maternal education | Primary school | 6(12.2%) | 36(73.5%) | 7(14.3%) | 0.001* | 3(6.1%) | 46(93.9%) | 0(0.0%) | 0.03* |
| | Junior high school | 18(9.9%) | 142(78.5%) | 21(11.6%) | | 9(5.0%) | 169(93.4%) | 3(1.7%) | |
| | High school | 12(9.0%) | 109(82.0%) | 12(9.0%) | | 9(6.8%) | 118(88.7%) | 6(4.5%) | |
| | Graduate | 2(10.0%) | 9(45.0%) | 9(45.0%) | | 0(0.0%) | 20(100.0%) | 0(0.0%) | |
| Economic status | Good | 8(14.5%) | 38(69.1%) | 9(16.4%) | 0.53 | 6(10.9%) | 49(89.1%) | 0(0.0%) | 0.07 |
| | Moderate | 27(8.9%) | 238(78.5%) | 38(12.5%) | | 12(4.0%) | 282(93.1%) | 9(3.0%) | |
| | Poor | 3(12.0%) | 20(80.0%) | 2(8.0%) | | 3(12.0%) | 22(80.0%) | 0(0.0%) | |
| Place of residence | Urban | 28(10.1%) | 210(75.8%) | 39(14.1%) | 0.44 | 14(5.1%) | 257(92.8%) | 6(2.2%) | 0.77 |
| | Rural | 10(9.4%) | 86(81.1%) | 10(9.4%) | | 7(6.6%) | 96(90.6%) | 3(2.8%) | |
| New born gender | Male | 18(9.3%) | 157(81.3%) | 18(9.3%) | 0.07 | 13(6.7%) | 171(88.6%) | 9(4.7%) | 0.99 |
| | Female | 20(10.5%) | 139(73.2%) | 31(16.3%) | | 8(4.2%) | 182(95.8%) | 0(0.0%) | |

cm: centimetre

on the results of chi-squared test (Table 5).

Discussion

Anaemia in pregnancy can result in the PTL, low birth weight, intrauterine growth retardation, increased risk of neonatal mortality, low physical activity capacity, immunodeficiency, and heart failure in mother (20, 21). Various mechanisms have been suggested regarding the effects of anaemia on pregnancy outcomes (17, 18). Among these mechanisms, two of them are of greater importance, including 1) the consequences of fetal low oxygenation due to low levels of haemoglobin, which result in intrauterine growth restriction, and 2) reduced size and surface of the placenta that causes low birth weight, as well as the presence of accompanying nutritional deficiencies in addition to anaemia (17, 18).

This study revealed a significant relation between Hb with low birth weight and the PTL in all three trimesters. In the present study, the relationship between maternal Hb and HCT levels with pregnancy outcomes were more

prominent in the first trimester. These findings were in line with the findings of the previously mentioned studies (20, 22-24). In a study carried on 903 Iranian pregnant women, low Hb and HCT levels were identified as predictors for low birth weight and the PTL (16). In a study carried out by Scholl et al. (2002), the optimum maternal HCT level related with the least possibility of the PTL was reported as 41%-44% (25).

The results of this study revealed no significant relation between high maternal Hb and HCT levels with pregnancy complications. It was in line with the findings of a previous study conducted on 609 pregnant women in Babol, Iran, which identified no relationship between maternal HCT level more than 40% and neonatal outcomes (26). Similarly, other studies have identified no significant relationship between high maternal HCT levels and pregnancy outcomes (27-29). In contrast, previous studies identified a significant relation between high maternal HCT levels and neonatal complications (23, 27).

High maternal HCT is known to be related to

the high volume of red blood cells and non-proportional increase in plasma volume, which is necessary for normal intrauterine growth; therefore, high HCT levels may result in negative pregnancy outcomes (30, 31). Moreover, according to the results of the present study, it was revealed that among the other parameters, only low maternal educational level and higher number of pregnancies were significantly associated with both the LBW and PTL. It was in line with the findings of the previous studies, which reported poor socioeconomic status, multiple pregnancy, and low maternal literacy as predictors of the LBW and PTL (7-9).

Conclusion

In this study, a possible reason for the lack of relationship between high maternal HCT level and pregnancy outcomes might be due to the limited number of the subjects with high HCT level, which can be considered as a limitation of this study. It is recommended that further studies should be performed on larger number of pregnant women with high HCT levels in order to evaluate the relationship between high maternal HCT levels and pregnancy outcomes.

The large sample size, which could represent the general population of pregnant women, can be considered as strength for this study. Furthermore, the present study included various confounders as neonatal weight and preterm labour in the analysis. On the other hand, since the aim of the study was to evaluate the relationship between maternal HB and HCT levels with neonatal weight and preterm labour, there could be no control over the number of women in each group. Further studies are required to assess this relationship among equal groups of the HCT and Hb levels using cluster sampling method.

Ethical Considerations

The present study was approved by the ethical committee of Sabzevar University of Medical Sciences and all the subjects signed informed consent forms prior to participation in the study.

Acknowledgements

The authors would like to thank the hospital authorities, nurses, and patients that efficiently

helped with the data collection.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Kramer MS, Barros FC, Demissie K, Liu S, Kiely J, Joseph KS. Does reducing infant mortality depend on preventing low birthweight? An analysis of temporal trends in the Americas. *Paediatric and Perinatal Epidemiology*. 2005; 19(6):445-451.
2. Alexander GR, Wingate MS, Mor J, Boulet S. Birth outcomes of Asian-Indian-Americans. *International Journal of Gynecology & Obstetrics*. 2007; 97(3): 215-220.
3. Karimian S, Mollamohammadi M, Jandaghi GR. Prevalence of low birth weight infants and its related factors in Qom delivery units, 2000. *Feyz Journals of Kashan University of Medical Sciences*. 2003; 7(3):76-80. (Persian)
4. Mansour E, Eissa A, Nofal LM, Kharboush I, Reda AA. Morbidity and mortality of low-birth-weight infants in Egypt. *Eastern Mediterranean Health Journal*. 2005; 11(4):723-731.
5. Delaram M, Akbari N. Weight gain in pregnancy and its correlation with birth weight of infants. *Knowledge & Health*. 2008; 3(2):39-43.
6. Bisai S, Mahalanabis D, Sen A, Bose K, Datta N. Maternal early second trimester pregnancy weight in relation to birth outcome among Bengalee Hindus of Kolkata, India. *Annals of Human Biology*. 2007; 34(1):91-101.
7. Lone FW, Qureshi RN, Emanuel F. Maternal anaemia and its impact on perinatal outcome. *Tropical Medicine & International Health*. 2004; 9(4):486-490.
8. Raqib R, Alam DS, Sarker P, Ahmad SM, Ara G, Yunus M, et al. Low birth weight is associated with altered immune function in rural Bangladeshi children: a birth cohort study. *The American Journal of Clinical Nutrition*. 2007; 85(3):845-852.
9. Kaneshi T, Yoshida T, Ohshiro T, Nagasaki H, Asato Y, Ohta T. Birthweight and risk factors for cardiovascular diseases in Japanese schoolchildren. *Pediatrics International*. 2007; 49(2):138-143.
10. FalahYakhdani M. The relationship between biological cognitive and psychosocial characteristics of parents and the weight of infant at the time of birth in Yazd Province. Department of exceptional children education, school of psychology Tehran: Tehran Azad Islamic University of Sciences and Researches; 2008. P. 85-100.
11. Garshasbi A, Fallah N. Maternal hematocrite level and risk of low birth weight and preterm delivery. *Tehran University Medical Journal*. 2006; 64(4):87-93.

12. Moutquin JM. Socio-economic and psychosocial factors in the management and prevention of preterm labour. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2003; 110(Suppl 20):56-60.
13. Cunningham F, Leveno K, Bloom S, Spong CY, Dashe J. *Williams obstetrics*. 24th ed. New York: McGraw-Hill; 2014.
14. Ingelfinger JR. Prematurity and the legacy of intrauterine stress. *New England Journal of Medicine*. 2007; 356(20):2093-2095.
15. Brabin BJ, Ginny M, Sapau J, Galme K, Paino J. Consequences of maternal anaemia on outcome of pregnancy in a malaria endemic area in Papua New Guinea. *Annals of Tropical Medicine and Parasitology*. 1990; 84(1):11-24.
16. Hajian KO, Asnafi N. The relationship between maternal hemoglobin. And hematocrit levels with neonatal pregnancy complication at birth. *Daneshvar Medicine*. 2006; 13(64):33-38. (Persian)
17. Idowu OA, Mafiana CF, Dopu S. Anaemia in pregnancy: a survey of pregnant women in Abeokuta, Nigeria. *African Health Sciences*. 2007; 5(4):295-299.
18. Ali EY, Adam GK, Ahmed S, Ali NI, Adam I. Maternal and neonatal hormonal profiles in anaemic pregnant women of eastern Sudan. *Journal of Obstetrics and Gynaecology*. 2009; 29(4):311-314.
19. Ren A, Wang J, Ye R, Li S, Liu J, Li Z. Low first-trimester hemoglobin and low birth weight, preterm birth and small for gestational age newborns. *International Journal of Gynecology & Obstetrics*. 2007; 98(2):124-128.
20. Lee HS, Kim MS, Kim MH, Kim YJ, Kim WY. Iron status and its association with pregnancy outcome in Korean pregnant women. *European Journal of Clinical Nutrition*. 2006; 60(9):1130-1135.
21. Bondevik G, Eskeland B, Ulvik RJ, Ulstein M, Lie R, Schneede J, et al. Anaemia in pregnancy: possible causes and risk factors in Nepali women. *European Journal of Clinical Nutrition*. 2000; 54(1):3-8.
22. Herberg S, Preziosi P, Galan P. Iron deficiency in Europe. *Public Health Nutrition*. 2001; 4(2b):537-545.
23. Cunningham FG, Leveno K, Bloom S, Hauth J, Gilstrap L, Wenstorm K. *Diseases and injuries of the fetus and newborn*. *Williams obstetrics*. 22nd ed. New York, NY: McGraw-Hill; 2005. P. 685.
24. Duthie SJ, King PA, To WK, Lopes A, Ma HK. A case controlled study of pregnancy complicated by severe maternal anaemia. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 1991; 31(2):125-127.
25. Rusia US, Madan NI, Agarwal NE, Sikka ME, Sood SK. Effect of maternal iron deficiency anaemia on foetal outcome. *Indian Journal of Pathology and Microbiology*. 1995; 38(3):273-279.
26. Nasiri Amiri F, Hajiahmadi M, Basirat Z. Maternal hematocrit status affecting pregnancy outcome. *Iranian Red Crescent Medical Journal*. 2007; 9(2):104-108.
27. Blankson ML, Goldenberg RL, Cutter G, Cliver SP. The relationship between maternal hematocrit and pregnancy outcome: black-white differences. *Journal of the National Medical Association*. 1993; 85(2):130.
28. Klebanoff MA, Shiono PH, Selby JV, Trachtenberg AI, Graubard BI. Anemia and spontaneous preterm birth. *American Journal of Obstetrics and Gynecology*. 1991; 164(1):59-63.
29. Siega-Riz AM, Adair LS, Hobel CJ. Maternal hematologic changes during pregnancy and the effect of iron status on preterm delivery in a West Los Angeles population. *American Journal of Perinatology*. 1997; 15(9):515-522.
30. Yildiz Y, Özgü E, Unlu SB, Salman B, Eyi EG. The relationship between third trimester maternal hemoglobin and birth weight/length; results from the tertiary center in Turkey. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2014; 27(7):729-732.
31. Jwa SC, Fujiwara T, Yamanobe Y, Kozuka K, Sago H. Changes in maternal hemoglobin during pregnancy and birth outcomes. *BMC Pregnancy and Childbirth*. 2015; 15(1):80.