

## Horseshoe-shaped Placenta and Preterm Labour: A Case Report

Hamideh Jafari (MSc)<sup>1</sup>, Robab Latifnejad Roudsari (PhD)<sup>2</sup>

<sup>1</sup> Lecturer, Department of Midwifery, School of Nursing and Midwifery, Torbat Heydarieh University of Medical Sciences, Torbat Heydarieh, Iran

<sup>2</sup> Associate Professor, Evidence-Based Care Research Centre, Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO	ABSTRACT
<i>Article type:</i> Case report	Horseshoe placenta is a variant of ring-shaped placenta. Ring-shaped placenta is a rare anomaly seen in fewer than 1 in 6000 deliveries, which could be seen as horseshoe shape due to atrophy in a portion of the ring. This report refers to a possible relationship between the horseshoe shape placenta and preterm labor; although it has not been yet validated in the literature. In this case report we present the placenta of a 35 years-old G2L1 woman who was referred to delivery unit of Shohada Hospital in Ghouch, Iran with starting of labor pain at 34 weeks of gestation who gave birth to a live-born female baby with 6/8 APGAR score. The amniotic fluid was meconium-stained. The patient had no evidence of fever, bleeding, infection. It is concluded that due to the risks to the fetus and mother caused by horseshoe placenta, its timely diagnosis during pregnancy using ultrasound is important and leads to early detection and better management of pregnancies complicated with placental abnormalities.
<i>Article History:</i> Received: 23-Nov-2013 Accepted: 1-Mar-2014	
<i>Key words:</i> Horseshoe-shaped placenta Placental abnormality Preterm labour Ring-shaped placenta	

► Please cite this paper as:

Jafari H, Latifnejad Roudsari R. Horseshoe-shaped Placenta and Preterm Labour: A Case Report. Journal of Midwifery and Reproductive Health. 2014;2(2):147-149.

### Introduction

Fetal development is a complex process, which depends on a variety of growth factors such as availability of nutrients and oxygen for the fetus. The placenta appears as the source of oxygen and nutrients for the fetus, therefore, it plays a central role in fetal growth. Placental processes depend on the size, morphology, blood flow, and transporter abundance of the placenta (1-4).

In terms of placental morphology, the shape of the chorionic plate of the human placenta is commonly round, with the umbilical cord roughly inserted at the center (4, 5). However, Salafia (2010) states that very few placentas appear truly round with a centrally inserted umbilical cord.

The placental shape has been commonly described as round, oval, bi- or multi-lobate, or otherwise irregular (6). Benirschke suggests that the mechanism by which a non-round placenta develops is unclear. However, the

findings support the notion of a secondary conversion from more normal placentation (7).

In less than 1 in 6,000 deliveries, the placenta is annular in shape, and sometimes a complete ring of the placental tissue is present. However, due to tissue atrophy in a portion of the ring, a horseshoe-shaped placenta becomes more common (Figures 1 & 2) (5).

Several studies have shown that abnormal placental growth is associated with adverse gestational outcomes, and could result in an impaired delivery of oxygen and nutrients to the fetus, thus leading to an abnormal fetal development (8, 11).

Horseshoe-shaped placenta, as a variant of ring-shaped placenta, appears to be associated with a greater likelihood of ante- and post-partum bleeding and fetal growth restriction (5).

This shape of placenta is so rare that there is no sufficient evidence to explain its

\* Corresponding author: Robab Latifnejad Roudsari, Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +98 511 8598016; Email: LatifnejadR@mums.ac.ir; rlatifnejad@yahoo.com



**Figure 1.** Fetus surface of a horseshoe shape placenta



**Figure 2.** Maternal surface of a horseshoe shape placenta

correlation with adverse gestational outcomes. Therefore, considering the importance of this type of placenta, due to its association with some fetal and maternal complications, in this report, we aimed to address a rare case of horseshoe-shaped placenta.

### Case Report

Herein, we report a horseshoe-shaped placenta with a non-central umbilical cord insertion (marginal cord). The subject was a 35 year-old gravida 2, para 1 woman, who was referred to the labour and delivery unit of Shohada Hospital in Quchan, Iran.

The subject had labor pain at 34 weeks' gestation and gave birth to a liveborn female infant, with cephalic presentation after two hours of labor, followed by a preterm vaginal delivery. The amniotic fluid was meconium-stained, and one- and five-minute APGAR scores were 6 and 8, respectively.

On first vaginal examination, the cervix was 70% effaced and 4 cm dilated, and the patient had no signs of fever, bleeding, infection, or history of medical complications throughout or before the pregnancy. The course of pregnancy was uneventful, and the subject received antenatal care in an outpatient unit.

### Discussion

In this report, by considering a possible relationship between the horseshoe-shaped placenta and preterm labor, an association between placental abnormality and other unidentified consequences can be suggested; however, the presence of such association has not been confirmed in the literature review.

Unfortunately, placental pathology is rarely included in the training of either obstetricians or

pathologists. As a result, the potential benefits of routine placental examination are in question. Salafia (1990) states that these benefits include the clarification of the causes of many adverse gestational outcomes, the improvement of future pregnancy risk assessments, and ascertainment of neonatal risk factors for long-term neurodevelopmental sequelae (12).

Penfold et al. (1979) demonstrated that there was no evidence regarding the association between fetal growth (its condition at birth) and cord eccentricity, placental shape and thickness, or the dimensions of the cord (13).

On the other hand, Yampolsky et al. (2009) showed that placental proportions, such as deviations of the placental shape from round and the relative thickness of the placenta modify placental functional efficiency, and in particular, explain some possible birth weight ranges in a normal population delivered from similar placental weights (4). Additionally, they indicated that the placenta with a displaced cord shows a markedly reduced transport efficiency, reflected in a larger value of  $\beta$ -HCG, and hence a smaller birth weight for a given placental weight (4).

Similarly, in our case, the displaced cord and placental shape might be caused by preterm labor due to reduced transport efficiency. In addition, according to Alwasel's findings (2013), tissue along the breadth of the placental surface may be more important than the tissue along the length in the transfer of nutrients from the mother to the fetus. In our case, the breadth of the placental surface was less than its length.

Costa et al. (2008) indicated that the determination of uterine artery Doppler and placental morphology (its shape and texture)

with ultrasound at 18-23 weeks of gestation could predict adverse perinatal outcomes. In their study, the placental shape was considered abnormal when depicting the following characteristics: placental length less than 10 cm, thickness more than 4 cm, or the length/thickness ratio less than 2.0 during the second trimester (15).

Therefore, due to the risks associated with horseshoe-shaped placenta for both the fetus and the mother, timely diagnosis during pregnancy (using ultrasound) is essential, and can lead to the early detection and proper management of pregnancies complicated with placental abnormalities.

### Conclusion

This report presented a possible relationship between the horseshoe-shaped placenta and preterm labor.

### Conflict of Interest

No conflict of interest exists.

### References

1. Mayeur S, Lukaszewski MA, Breton CH, Storme L, Vieau D, Lesage J. Do neurotrophins regulate the fetoplacental development? *Medical Hypotheses*. 2011; 76(5):726-728.
2. Li D, Wi S. Maternal Placental Abnormality and the Risk of Sudden Infant Death Syndrome. *American Journal of Epidemiology*. 1999; 149(7):608-611.
3. Jansson T, Powell TL. Role of the placenta in fetal programming: underlying mechanisms and potential interventional approaches. *Clinical Science*. 2007; 113:1-13.
4. Yampolsky M, Salafia CM, Shlakhter O, Haas D, Eucker B, Thorp J. Centrality of the Umbilical Cord Insertion in a Human Placenta Influences the Placental Efficiency. *Placenta*. 2009; 30(12): 1058-1064.
5. Cunningham FG, Leveno KJ, Bloom SL, Hauth JC, Rouse DJ, Spong CY. *Williams obstetrics*. 23rd ed. New York: McGraw-Hill; 2010.
6. Salafia CM, Yampolsky M, Misra DP, Shlakhter O, Haas D, Eucker B, et al. Placental surface shape, function, and effects of maternal and fetal vascular pathology. *Placenta*. 2010; 31:958-962.
7. Benirschke K, Kaufmann P, Baergen R. Architecture of normal villous trees, chapter 7. In: *Pathology of the human placenta*. 5th ed. New York: Springer Verlag; 2006. p. 121-159.
8. Kingdom J, Huppertz B, Seaward G, Kaufmann P. Development of the placental villous tree and its consequences for fetal growth. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2000; 92:35-43.
9. Naeye RL. Causes and consequences of placental growth retardation. *JAMA*. 1978; 239:1145-1147.
10. Wallace JM, Aitken RP, Milne JS, Hay Jnr WW. Nutritionally-Mediated Placental Growth Restriction in the Growing Adolescent: Consequences for the Fetus. *Biology of Reproduction*. 2004; 71:1055-1062.
11. Baptiste-Roberts K, Salafia CM, Nicholson WK, Duggan A, Wang NY, Brancati FL. Maternal risk factors for abnormal placental growth: The national collaborative perinatal project Kesha. *BMC Pregnancy and Childbirth*. 2008;8(1): 1-7.
12. Salafia CM, Vintzileos AM. Why all placentas should be examined by a pathologist in 1990. *American Journal of Obstetrics and Gynecology*. 1990;163:1282-1293.
13. Penfold P, Drury L, Lewis L, Royston JP, Hytten FE. Case note descriptions of the placenta: are they worthwhile? *BJOG: An International Journal of Obstetrics & Gynaecology*. 1979; 86(5): 337-339.
14. Alwasel SH, Harrath AH, Aljarallah JS, Abotalib Z, Osmond C, Al Omar SY, et al. The velocity of fetal growth is associated with the breadth of the placental surface, but not with the length. *American Journal of Human Biology*. 2013; 25(4):534-537.
15. Costa SL, Proctor L., Dodd JM, Toal M, Okun N, Johnson JA, et al. Screening for Placental Insufficiency in High-risk Pregnancies: Is Earlier Better? *Placenta*. 2008; 29(12): 1034-1040.