

The Relationship between Maternal-Fetal Attachment and Mother-Infant Attachment Behaviors in Primiparous Women Referring to Mashhad Health Care Centers

Mahin Taffazoli (MSc)¹, Maryam Montakhab Asadi (MSc)^{2*}, Seyed Amir Aminyazdi (PhD)³, Mohamad Taghi Shakeri (PhD)⁴

¹ Lecture, Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran

² MSc in Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran

³ Associate Professor, Department of Psychology, School of Psychology and Educational Sciences, Ferdowsi University, Mashhad, Iran

⁴ Professor of Biostatistics, Department of Social Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article type:
Original article

Article History:
Received: 7-Aug-2014
Accepted: 21-Jan-2015

Key words:
Maternal-fetal attachment
Mother-infant attachment behaviors
Primiparous
Women

ABSTRACT

Background & aim: Mother-infant bonding and interactions after childbirth are shaped by maternal-fetal attachment during pregnancy. Although many studies have shown the positive correlation between maternal-fetal attachment and mother-infant attachment behaviors, some controversial studies have shown otherwise. Therefore, this study aimed to evaluate the correlation between maternal-fetal attachment and mother-infant attachment behaviors in primiparous women.

Methods: This descriptive correlational study was conducted on 100 primiparous women, referring to the selected health care centers of Mashhad. Data were collected using Cranley's maternal-fetal attachment scale, Avant's mother-infant attachment tool, Edinburgh postnatal depression scale, and a demographic/obstetric questionnaire including demographic data, obstetric information, delivery outcomes, and postpartum data. Pregnant women with a gestational age of 35-41 weeks, who met the inclusion criteria, completed Cranley's questionnaire, as well as the demographic/obstetric questionnaire. Four and eight weeks after delivery, the subjects were asked to complete the Edinburgh questionnaire and postpartum information; then, they were asked to breastfeed their infants on a chair in a quiet place for 15 minutes. The researcher observed the mothers' behaviors toward their neonates. For data analysis, descriptive and analytical tests were performed, using SPSS version 16.

Results: There was a direct positive relationship between maternal-fetal attachment and mothers' emotional behaviors toward infants four and eight weeks after delivery. However, four and eight weeks after childbirth, no significant correlation was found between maternal-fetal attachment and mothers' caring behaviors.

Conclusion: According to the findings, maternal-fetal attachment is one of the most important factors for mother-infant attachment. These findings could be applied for enriching mother-infant attachment behaviors during pregnancy.

► Please cite this paper as:

Taffazoli M, Montakhab Asadi M, Aminyazdi SA, Shakeri MT. The Relationship between Maternal-Fetal Attachment and Mother-Infant Attachment Behaviors in Primiparous Women Referring to Mashhad Health Care Centers. Journal of Midwifery and Reproductive Health. 2015; 3(2): 318-327.

Introduction

Infancy is one of the most important periods in an individual's life in terms of growth and development. Mental health of a neonate is as important as his/her physical health (1-3). Mothers show their emotional attachment to their infants by behaviors such as smiling,

calling their names, attentive responses to their movements, touching, hugging, direct eye contact, and kissing (4).

Mother-infant attachment reflects the quality of mother's emotional feelings and behaviors toward her baby; this bonding emerges in

* Corresponding author: Maryam Montakhab Asadi, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran. E-mail: MontakhabM891@mums.ac.ir

behaviors which indicate the mother's attention and care. These behaviors include affectionate, proximity, and caring behaviors (5).

According to previous studies, women with stronger attachment to their children are most often sensitive to their children's needs. This sensitivity positively influences the infant's future psychosocial function and development of personality traits such as curiosity, sociability, self-confidence, independence, collaboration, and honesty, which are gradually formed (6).

On the other hand, children with insecure mother-infant bonding experience less emotional and mental development, poor social interactions, and stronger tendencies to escape from school; moreover, they are less able to form long-term relations and show more aggressive behaviors, compared to others (7, 8).

Several factors affect maternal-fetal attachment such as mother's personality disorders including psychiatric diseases, depression, bipolar disorders, and schizophrenia (9), marital satisfaction, instable family status (10), mother's early attachment experiences during childhood, and support provided by others during and after pregnancy (11, 12).

According to previous research, mother-infant attachment and interactive behaviors are formed during pregnancy by maternal-fetal attachment (13-15). Speckhard believes that maternal-fetal attachment is formed before the neonate's birth when the mother begins to form a mental image of the baby and imagines supporting her infant (16). Cranley also stated that changes in the quality of mother-infant relation starts before birth (13).

Maternal-fetal attachment refers to the relation between mother and fetus, which is related to the mother's mental image of the infant. This form of attachment is shaped during pregnancy, gradually improves in the third semester, and continues after delivery (17, 18). The emerging maternal-fetal attachment could be a predictor for the attitude and performance of mothers after delivery, as well as mother-infant interactions and attachment patterns after birth (19).

Mothers, who are sensitive about their maternal role during pregnancy, remain sensitive after childbirth (20). Pregnant women form an attachment to the fetus by talking to it,

touching the abdomen, and paying attention to its movements. These mothers are ready to form a pleasant communication with their babies after delivery (21).

Moreover, increasing maternal-fetal attachment leads to healthy eating, alcohol withdrawal, positive attitudes towards the fetus, attention to fetal movements, and other interactive behaviors with the fetus. All these behaviors could lead to the acceptance of maternal role and promotion of physical and psychological health in mothers and infants (22).

As Reading et al. indicated (1982), strong maternal-fetal attachment is positively associated with healthy behaviors in mothers (23); these healthy behaviors lead to the promotion of maternal, fetal, and neonatal health (24). As Leifer (1986) (25), Fuller (1989) (26), and Muller (1996) (27) reported in different studies, emotional behaviors and maternal-fetal attachment during pregnancy could be predictors of mother-infant attachment behaviors.

Although most studies have shown a positive relationship between maternal-fetal attachment and mother-infant attachment behaviors, some controversial studies have reported otherwise. For instance, Davis (1989) reported no significant difference in the attachment behaviors of mothers with and without abdominal touching during the third semester of pregnancy (28). Similarly, Cranley (1981) found no relationship between maternal-fetal attachment and maternal feelings in the first three days after birth (29).

Considering the results of these controversial studies, personal/cultural differences, and the importance of the advisory, supportive, and administrative role of midwives during and after pregnancy, this study aimed to determine the relationship between maternal-fetal attachment and mother-infant attachment behaviors. By determining the influential factors, applying effective strategies, and preventing the infants' deprivation of maternal affection, the psychological health of mothers and infants can be promoted.

Materials and Methods

This descriptive correlational study was conducted on 100 primiparous women, who

referred to the selected health care centers of Mashhad for pregnancy care. The sample size was calculated based on a pilot study on 20 primiparous women, using the following formula:

$$n = \left(\frac{Z_{1-\alpha/2} + Z_{1-\beta}}{\omega} \right)^2 + 3$$

Considering CI=95%, $\alpha=5\%$, and power=80%, β was estimated at 20%. Among different sample sizes and different variables in previous studies, the maximum sample size was related to the variable of "exclusive feeding duration in 8 weeks", which was a dependent variable in the study ($n=33$) ($r=0.47$). This number was multiplied by three due to the nature of the study (cluster and stratified sampling).

Considering the 10% dropout, 110 participants were introduced to the study. However, 10 subjects were eliminated due to the mother's lack of consent to continue the study, two because of neonatal hospitalization, two due to depression, three for incomplete daily reports of neonatal nutrition, and one due to suffering from severe nipple fissure. Eventually, 100 participants were evaluated.

Sampling was conducted in 3 steps: stratified sampling, clustering, and simple non-probable sampling from August to January 2013. In the first step, five health care centers were stratified, and based on the patient coverage of each cluster, some clusters were randomly selected by drawing lots. According to the frequency of referrals for prenatal care, a quota was set for each cluster. Days of hospital visit were determined by drawing lots. On these days, based on the number of covered patients in each center, eligible participants were selected.

The inclusion criteria were as follows: 1) Iranian nationality; 2) residing in Mashhad; 3) 18-35 years of age; 4) literacy; 5) no prior history of medical diseases; 6) no prior history of drug addiction, hookah smoking, or cigarette smoking; 7) no history of infertility; 8) gestational age of 25-41 weeks; 9) primiparous singleton pregnancy; 10) low-risk pregnancy; 11) neonatal weight appropriate for gestational age (2500-4000 g); and 12) not studying medical sciences.

The exclusion criteria were as follows: 1) unwillingness to continue the study; 2) taking psychiatric medications; 3) major stress in the last 6 months; 4) postpartum complications; 5) neonatal hospitalization or separation of the neonate from the mother; 6) stressful events during the study period; 7) unavailability of the participants four and eight weeks after delivery (or two days after the set dates); and 8) depression after delivery.

Ethical considerations were observed throughout the study and an approval letter was obtained from the ethics committee of Mashhad University of Medical Sciences (MUMS). In addition, an introduction letter was obtained from the School of Nursing and Midwifery for presentation to the authorities of health care centers. Participants, who met the inclusion criteria, were referred for pregnancy care at the gestational age of 35-41 weeks.

Informed consents were obtained from the participants and questionnaires of demographic data, obstetric information, and maternal-fetal attachment were handed to them. They answered the questionnaires at the presence of the researcher and the researcher answered their questions. The participants and researcher exchanged their phone numbers so that they could contact each other about the time of delivery. The researcher contacted the participants before the delivery to ensure the exact time.

In the next step, the researcher, knowing the time of delivery, excluded the subjects in case they met the mentioned criteria; otherwise, the questionnaire of obstetric and neonatal data was completed via phone calls, and the participants were reminded to visit the hospital within four and eight weeks after delivery.

During mothers' visits after four and eight weeks, Edinburgh postnatal depression scale was completed. Then, they were asked to sit in a calm and quiet place to breastfeed their infants and play with them for 15 min. The researcher observed and recorded mothers' behaviors toward their infants. Each minute was divided to two 30 seconds; in the first 30 seconds, the behaviors were observed and during the second half, the behaviors were recorded.

Data collection tools

Selection form

This form, completed via interview, consisted of the inclusion and exclusion criteria, which were based on the research objectives, latest published articles, and consultation with faculty members.

Demographic and obstetric questionnaire

This questionnaire consisted of 2 sections. The first part included 9 demographic questions which were completed at the beginning of the study. The second section included 12 questions related to the current pregnancy status of mothers; ten items were answered via interview and two items were completed using examination or mothers' medical records.

Cranley's maternal-fetal attachment questionnaire

This questionnaire consisted of 24 statements, graded by 5-point Likert scale: strongly agree (5), agree (4), not sure (3), disagree (2), and strongly disagree (1). Question No. 22 was reverse scored. The minimum and maximum scores were 24 and 120, respectively, and higher scores indicated stronger attachment.

Delivery and neonatal questionnaire

This questionnaire included two sections. The first part had three questions related to the time of delivery, maternal satisfaction, and childbirth experience and was completed via phone calls. The second part consisted of 10 questions related to neonatal gender and weight and parental satisfaction with the baby's gender.

Avant's questionnaire of mother-infant attachment behaviors

This questionnaire included three groups of behaviors: emotional behaviors, proximity behaviors, and caring behaviors, observed for 15 min. In the first 30 seconds of each minute, the behaviors were observed and during the second half, these behaviors were recorded; each observed behavior was recorded once per minute.

In total, the observation took 15 min; therefore, each behavior was observed for a maximum of 15 times in 15 minutes. In total, 11 behaviors could be observed in 15 min and each

observed behavior in one minute was allocated one score (total score=165). The sum of scores presented mother-infant attachment and higher scores indicated stronger attachment.

Edinburgh postnatal depression scale

This questionnaire consisted of 10 short phrases. Each answer was given a score of 0 to 3, based on the severity of the symptoms. Questions No. 1, 2, and 4 were scored from 0 to 3 and the rest were scored from 3 to 0. The total score was calculated by summing the scores. A total score of ≥ 10 indicated depression and a total score of < 10 showed the absence of depression. This questionnaire was completed 4 and 8 weeks after delivery by the mothers.

In order to confirm the validity of demographic and obstetric questionnaire, as well as the delivery and neonatal questionnaire, content validity was applied. At first, the questionnaires were prepared, based on the latest published papers, and then, 10 faculty members were asked to comment on the questionnaires. The questionnaires were revised based on the given comments and the final version was prepared for the study.

The validity of Avant's questionnaire was first determined by Avant (30). This questionnaire was translated to Farsi and by applying some changes in the questionnaire, its validity was confirmed. The validity of this questionnaire was also approved by Tousi et al. (2011) (32). The reliability of this scale was also confirmed via inter-rater reliability by Vakilian et al. (1998) (31).

Cranley's maternal-fetal attachment questionnaire was first applied by Cranley (1980) and its validity was confirmed (33). This scale was translated to Farsi by Khorramrudi (2001) in Iran and its validity was confirmed using content validity (34). Abbasi et al. also confirmed the validity of the Farsi version using content validity (2010) (35).

In addition, the reliability of this questionnaire was first approved by Cranley (1981) (33). The reliability of the Farsi version was confirmed by Khorramrudi (2001) via test-retest ($r=0.85$) (34) and by Abbasi et al. (2010) by calculating Cronbach's alpha ($\alpha=0.8$) (35). In the present study, Cronbach's alpha was calculated to be 0.08. Moreover, the reliability of

Edinburgh depression scale has been confirmed in Iran in several studies such as those by Namazi (1994) ($r=0.9$) (36) and Montazeri et al. (2007) (37).

For data analysis, Chi-square, Mann-Whitney U test, and Spearman's correlation coefficient were performed, using SPSS version 16. Moreover, descriptive statistics, absolute and

relative frequency, and mean and standard deviation were calculated.

Results

The subjects were within the age range of 18-35 years (26 ± 4.67 years) and gestational age of 36-41 weeks (37 ± 1.28) (Table 1). In total, 77 women (77%) were housewives and 43 (43%)

Table 1. Demographic characteristics of participants in the study

Variables	Sub-groups	Number (percentage)
Maternal education Number (percentage)	Minimum literacy (reading and writing)	10 (10.0)
	Primary level education	20 (20.0)
	Junior high education	27 (27.0)
	High school diploma	30 (30.0)
	Academic education	13 (13.0)
Total 100		
Husband's educational level Number (percentage)	Minimum literacy (reading and writing)	4 (4.0)
	Primary level education	27 (27.0)
	Junior high education	27 (27.0)
	High school diploma	33 (33.0)
	Academic education	9 (9.0)
Total 100		
Maternal occupation Number (percentage)	Housewife	77 (77.0)
	Employed	13 (13.0)
	Student	10 (10.0)
Total 100		
Husband's occupational status Number (percentage)	Employed	15 (15.5)
	Worker	82 (84.5)
Total 100		
Breastfeeding training after delivery Number (percentage)	Yes	74 (74.0)
	No	26 (26.0)
Total 100		
Mode of delivery Number (percentage)	Normal vaginal delivery	56 (56.0)
	Emergency cesarean section	38 (38.0)
	Elective cesarean section	6 (6.0)
Total 100		
First lactation after delivery Number (percentage)	Immediately after delivery	54 (54.0)
	Less than 2 hours	41 (41.0)
	More than 2 hours	5 (5.0)
Total 100		
First skin contact after delivery Number (percentage)	Yes	63 (63.0)
	No	37 (37.0)
Total 100		
Marital satisfaction Number (percentage)	Not at all	0 (0.0)
	Low	1 (1.0)
	Relatively low	15 (15.0)
	Relatively high	45 (45.0)
	High	39 (39.0)
	Very high	0 (0.0)
Total 100		
Interest in children Number (percentage)	Not at all	75 (75.0)
	Medium	9 (9.0)
	High	16 (16.0)
Total 100		

Table 2. Demographic characteristics of participants in the study

Variables	SD±Mean
Maternal age	67.26±4
Gestational age	28.37±1
Neonatal weight	67.18±456.3524
BMI	85.22±1.25

had high school diploma (or higher) (Table 1). Mean body mass index (BMI) before pregnancy

was 25.22±1.85 kg/m². As the results showed, 56% of participants had normal delivery (Tables 1 & 2). Other demographic data are presented in Tables 1 & 2. Correlation of descriptive characteristics associated with total score of mother-infant attachment behaviors in 4 and 8 weeks after birth are presented in Tables 3 & 4 and results of mother-infant attachment behavior in the influential variables are

Table 3. Correlation coefficients for descriptive characteristics associated with the total score of mother-infant attachment behaviors four and eight weeks after birth

Variables	Mother-infant attachment behaviors after 4 weeks		Mother-infant attachment behaviors after 8 weeks	
Maternal age	Spearman	P=0.106 r=0.336	Spearman	P=0.247 r=0.135
Maternal education	Spearman	P< 0.001	Spearman	P< 0.001
Husband's education	Spearman	P=0.474	Spearman	P=0.491
Marital satisfaction	Spearman	P< 0.001	Spearman	P< 0.001
Gestational age	Spearman	P=0.211 r=-0.170	Spearman	P=0.300 r=-0.225
BMI	Spearman	P=0.478 r=-0.322	Spearman	P=0.465 r=-0.155
Mode of delivery	Chi square	P=0.438	Chi square	P=0.430
Neonatal weight	Spearman	P=0.170 r=0.178	Spearman	P=0.359 r=0.193
First lactation after delivery	Spearman	P< 0.001	Spearman	P< 0.001

Table 4. Correlation coefficients for descriptive characteristics associated with the total score of mother-infant attachment behaviors four and eight weeks after childbirth

Variables	Sub-groups	Mother-infant attachment behaviors 4 weeks after delivery (Mean±SD)		Mother-infant attachment behaviors 8 weeks after delivery (Mean±SD)	
Maternal occupational status	Housewife	1.25±0.43		1.26±0.44	
	Employed	1.28±0.48		1.28±0.48	
	Student	1.28±0.48		1.14±0.37	
F=0.267 P=0.066	F=0.027 P=0.212	Spearman		Results	
Husband's occupation	Employed	1.26±0.44		1.21±0.42	
	Worker	1.29±0.46		1.29±0.46	
F=1.547 P=0.464	F=1.193 P=0.433	Chi-square		Results	
First skin contact after delivery	Immediately after delivery	1.22±0.42		1.22±0.44	
	One hour after delivery	1.25±0.43		1.25±0.43	
	Two hours after delivery	1.55±0.52		1.33±0.50	
F= 1.628 P< 0.001	F= 1.628 P< 0.001	Chi-square		Results	
Breastfeeding training after delivery	Yes	1.25±0.43		1.30±0.46	
	No	1.26±0.44		1.20±0.40	
F= 1.524 P=0.506	F= 0.019 P=0.695	Chi-square		Results	
Interest in children	Not at all	1±0.00		1±0.00	
	Medium	1.15±0.37		1.26±0.45	
	High	1.29±0.46		1.29±0.46	
F= 0.213 P=0.004	F= 0.556 P=0.027	Chi-square		Results	

Table 5. The results of regression model based on mother-infant attachment behaviors in the presence of effective variables

Effective variables	Attachment four weeks after delivery (P-value)	Attachment eight weeks after delivery (P-value)
Maternal education	001.<0	05.>0
Marital satisfaction	05.>0	001.<0
Interest in children	020.0	004.0

presented in Table 5. The results of Spearman's correlation coefficient showed a direct positive association between maternal-fetal attachment and emotional and proximity behaviors in mother-infant bonding four and eight weeks after delivery ($P < 0.001$) (Table 6). In fact, stronger maternal-fetal attachment was associated with more emotional and proximity behaviors. However, no significant relationship was found between maternal-fetal attachment and mother-infant caring behaviors at week 4) (Table 6).

Discussion

Maternal-fetal attachment is an important part of pregnancy that facilitates future communication between the mother and neonate (38). More attached mothers are able to keep the fetus and neonate healthy, whereas less attached mothers experience various problems and are unable to maintain the health of their fetus or neonate (39).

The present study showed a significant direct relationship between maternal-fetal attachment and emotional and proximity behaviors; i.e., stronger maternal-fetal attachment was accompanied by more emotional and proximity behaviors in mother-infant relationship. In this regard, Abbasi et al. (2008) showed that increased maternal-fetal attachment leads to the improvement of mother's mental health and desirable communication between mother and neonate (35).

Nagata et al. (2003) also showed a significant correlation between intrauterine/postpartum attachment and attachment one year after delivery (40). In a study by Vakilian et al. (1386), emotional behaviors had the maximum score among other attachment behaviors (31). Findings of the present study were in agreement with studies by Davachi (2000) (4), Carter (1981) (38) and Griss (43). In other words, mothers who paid more attention to their fetus

Table 6. Correlation of maternal-fetal attachment with emotional, caring, and proximity behaviors four and eight weeks after childbirth

Variables	Four weeks after delivery Rs (P)	Eight weeks after delivery Rs (P)
Emotional behaviors	0.752(< 0.001)	0.670(< 0.001)
Proximity behaviors	0.753(< 0.001)	0.626(< 0.001)
Caring behaviors	0.072(0.478)	0.120(0.233)

showed more affection and care toward their babies after delivery; they smiled more often to their infants and hugged and kissed them more regularly.

Fuller in his study on maternal-fetal attachment stated that attachment before delivery is related to mother-infant attachment during the first days of birth (26). Griss concluded that attachment before delivery could predict attachment 4-6 weeks after delivery. Leifer (1986) (25) and Muller (1996) (27) in separate studies reported that maternal-fetal attachment during pregnancy could be considered a predictor for mother-infant attachment behaviors.

Although many studies have shown a positive relationship between maternal-fetal attachment and mother-infant attachment behaviors, some controversial results have been reported. For instance, Davis (1989) found no significant difference between attachment behaviors of mothers who touched their abdomen during the third semester of pregnancy and others (28).

Similarly, Cranley found no relationship between maternal-fetal attachment and maternal feelings within the first 3 days after birth (29). The reason behind this discrepancy could be the application of different instruments for observing mother-infant attachment behaviors and the short follow-up period, which was only 2-3 days after birth in these studies.

Mihara et al. (2005) studied attachment behaviors in Japanese and Brazilian populations and concluded that women's feelings are similar, although there are some differences in behaviors between these populations. For instance, Japanese mothers mostly looked at their babies without touching them, while Brazilian mothers looked at their babies less and spent more time on taking care of their

neonates. These behaviors were related to cultural and socioeconomic differences between these countries (44).

According to the findings of the present study, no significant correlation was found between maternal-fetal attachment and mother-infant caring behaviors. This result was in contrast with the findings by Brimnezhad et al. (2011) (45) and Vakilian et al. (2008) (31). In the mentioned studies, the minimum score was given to caring behaviors. According to the present research and other conducted studies, maternal-fetal attachment is one of the influential factors for mother-infant attachment behaviors. These findings could be applied for enriching mother-infant attachment behaviors during pregnancy.

Application of results

The results could be applied for future health care planning and policy making. For instance, consultation and educational programs can be held for health care providers to focus on maternal-fetal attachment and mother-infant bonding.

Limitations

Despite the researchers' effort to eliminate and control the confounding factors, some of them were out of control. First, the differences in participants' personalities and emotional status affected their answers to the questionnaires; this problem was relatively controlled by the random selection of health care centers.

Second, despite assuring participants about the confidentiality of the data and providing proper conditions for answering the questions, some participants might have answered the questions less precisely; this was also uncontrollable by the researcher. Finally, answering the questionnaires in a limited amount of time and breastfeeding out of the house at the presence of the researcher might have influenced the given answers.

Acknowledgements

This study was part of an M.Sc. research proposal, approved by MUMS (code: 911204). The authors would like to thank MUMS for the full financial support, the respected staff at

health care centers for their cooperation, and mothers for their participation in the study.

References

1. Carter JL. Promoting maternal attachment through prenatal intervention. *MCN: The American Journal of Maternal/Child Nursing*. 1981; 6:107-112.
2. Behrman RE, Kliegman RM, Jenson HB (eds). *Nelson Textbook of Pediatrics*. 17th ed. Philadelphia: Saunders. 2004; P: 157-161.
3. Schanler RJ. The use of human milk for premature infants. *Pediatric Clinics of North America*. 2001; 48(1):207-219.
4. Carter JL. Promoting maternal attachment through prenatal intervention. *MCN: The American Journal of Maternal/Child Nursing*. 1982; 16:107-112.
5. Virginia LC, Nancy L, Associates, Inc. *Infant attachment: what we know now*. U.S department of health and human services 1991. Available at: <http://aspe.hhs.gov/daltcp/reports/inatrpt.pdf>.
6. Mc dad E. *Psychological Preparation for Motherhood and its Association with Postpartum Outcomes* [PhD dissertation]. Eugene: University of Oregon; 2008.
7. Wilkerson ML. *Maternal Role Attainment Among First Time Mothers Of Full Term Infants And Premature Infants As A Function Of Planned VS. Unplanned Pregnancy* [PhD dissertation]. Texas: University of Houston; 1996.
8. Muller M. A questionnaire to measure mother to infant attachment. *Journal of Nursing Measurement*. 1994; 2(2):129-141.
9. Gelder M. *Oxford Textbook of psychiatry*. 3rd ed. Oxford: Oxford University Press; 1996.
10. Yarcheski A, Mahon NE, Yarcheski TJ, Hanks MM, Cannella BL. A meta-analytic study of predictors of maternal-fetal attachment. *International Journal of Nursing Studies*. 2009; 46(5):708-715.
11. Pillitteri A. *Maternal and child health nursing: Care of the Childbearing and Childrearing Family*. Philadelphia: Lippincott Williams & Wilkins; 2013. p: 601-616.
12. Cannella BL. Maternal -fetal attachment: an integrative review. *Journal of Advanced Nursing*. 2005; 50(1): 60-68.
13. Jose M, Graciela N, Liliana B, Ana M, Armando G. Maternal and perinatal factors influencing the duration of exclusive Breast feeding during the first 6 month of life. *Journal of Human Lactation*. 2003; 19(2):136-144
14. Losch M, Dungy CI, Russell D, Dusdieker LB. Impact of attitudes on maternal decisions

- regarding infant feeding. *The Journal of Pediatrics*. 1995; 126:507-514.
15. Salisbury A, Law K, LaGasse L, Lester B. Maternal-fetal attachment. *JAMA*. 2003; 289(13):1701.
 16. Speckhard A. Traumatic death in pregnancy: The significance of meaning and attachment. In Figley CR, Bride BE, Mazza N (eds), *Death and trauma: The traumatology of grieving*. Washington, DC: Taylor and Francis; 1997. p: 67-100.
 17. Pillite A. *Maternal and child health nursing*. 4th ed. UK: Lippincott; 2003; p: 201-226
 18. Lawson KL, Turriff-Jonasson SI. Maternal serum screening and psychosocial attachment to pregnancy. *Journal of Psychosomatic Research*. 2006; 60: 371-378.
 19. Foster SF, Slade P, Wilson K. Body image, maternal-fetal attachment and breast-feeding. *Journal of Psychosomatic Research*. 1996;41: 181-184
 20. Mikhail MS, Freda MC, Merkatz RB, Polizzotto R, Mazloom E, Merkatz IR. The effect of fetal movement counting on maternal attachment to fetus. *American Journal of Obstetrics and Gynecology*. 1991; 165(4):988-991.
 21. Eileen F R. *The Relationship between Prenatal Maternal Attachment, Postpartum Depressive Symptoms and Maternal Role Attainment* [PhD dissertation]. Chicago: University of Chicago; 1994.
 22. Reading AE, Campbell S, Cox DN, Sledmere CM. Health beliefs and health care behavior in pregnancy. *Obstetrical & Gynecological Survey*. 1983; 38(3):140-141.
 23. Lindgren K. Relationships among maternal-fetal attachment, prenatal depression and health practices in pregnancy. *Research in Nursing and Health*. 2001; 24: 203-217.
 24. Huang HC, Wang SY, Chen CH. Body Image, Maternal-Fetal Attachment, and Choice of Infant Feeding Method: A Study in Taiwan. *Birth*. 2004; 31(3):183-188.
 25. Leifer M. *Psychological effects of motherhood*. New York: Prager; 1980.
 26. Fuller JR. Early patterns of maternal attachment. *Health Care for Women International*. 1990; 11:433-446.
 27. Müller ME. Prenatal and postnatal attachment: A modest correlation. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 1996; 25(2): 161-166.
 28. Davis MS, Akridge KM. The effect of promoting intrauterine attachment in primiparas on postdelivery attachment. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 1987; 16(6): 430-437.
 29. Cranly MS. Development of a tool for the measurement of maternal attachment during pregnancy. *Nursing Research*. 1981; 30: 281-284.
 30. Tilokskulchai F, Phatthanasiriwethin S, Vichitsukon K, Serisathien Y. Attachment behaviors in mothers of premature infants: a descriptive study in Thai mothers. *The Journal of Perinatal & Neonatal Nursing*. 2002; 16(3):69-83.
 31. Vakilian K, Khatamidoost F, Khorsandi M. Effect of kangaroo mother care immediately after birth on maternal attachment behavior before hospital discharge. *Medical Journal of Hormozgan University*. 2007; 11(1):57-63.
 32. Toosi M, Akbarzadeh M, Zare N, Sharif F. Effect of Attachment Training on Anxiety and Attachment Behaviors of first-time Mothers. *Hayat Journal*. 2011; 17(3):69-79
 33. Siddiqui A, Hagglof B. Does maternal prenatal attachment predict postnatal mother-infant interaction?. *Early Human Development*. 2000; 59: 13-25.
 34. Khoramrody R. *The effect of mothers touch on Maternal Foetal Attachment* [PhD dissertation]. Tehran: Iran University of Medical Sciences; 2000.
 35. Abbasi A, Tafazoli M. The effect of education of maternal attachment behavior on maternal-fetal attachment in primiparus women. *Journal of Hamadan Nursing and Midwifery University*. 2009; 17(1): 35-40.
 36. Namazi SH. *The roll e of psychological and demographic variables on postpartum depression* [PhD dissertation]. Tehran: Iran University of Medical Sciences; 1993.
 37. Montazeri A, Torkan B, Omidvar S. The Edinburgh postnatal depression scale (EPDS): translation and validation study of the Iranian version. *BMC Psychiatry*. 2007; 7: 11.
 38. Andrade S, Atkins M, Battersy S, Buchanan P, Cowbrough K, Duncan J, et al. National Institute for Health and Clinical Excellence. A peer-support programme for woman who breastfeed commissioning guide implementing NICE guidance; 2008.
 39. Sandbrook SP, Adamson Macedo EN. Maternal-fetal attachment: searching for a new definition. *Neuro Endocrinology Letters*. 2004; 25(1): 169-182.
 40. Nagata M, Nagai Y, Sobajima H, Ando T, Honjo S. Depression in mother and maternal attachment results from a follow-up study at 1years postpartum. *Psychopathology*. 2002; 36: 142-151.
 41. Davachi A, Khoramroodi R, Shahpoorian F, Haghani H. Fetal touching! Does it improve maternal fetal, or meternal newborn attachment?. *Iran Journal of Nursing*. 2000; 13(24):16-24. [Persian]

42. Vakilian K. The effect of contact immediately after delivery between mother and infant on behavior attachment in nulliparous women before discharge, one and three month after birth in the teaching hospital in Tehran [PhD dissertation]. Tehran: Iran University of Medical Sciences; 1992.
43. Nematbakhsh F, Kordi M, Sahebi A, Esmaeeli H. The effect of mother – infant skin to skin contact on mother’s attachment. *The Quarterly Journal of Fundamentals of Mental Health*. 2007; 33(9): 25-32.
44. Cassano CA, Maehara, S. Japanies and Brazilian maternal bonding behavior to word preterm infant: A comparative study. *Journal of Neonatal Nursing*. 1998; 4(1): 23-2.
45. Barimnejad L, Mehrnoosh L, Seyedfatemi N, Haghani H. The effect of Empowerment Program on Mother-Infant Interaction and Weight Gain in Preterm Infants Hospitalized in Neonatal Intensive Care Unit. *Zahedan Journal of Research in Medical Sciences*. 2012; 14(9):12-17.