

Acute Burns during Pregnancy: A Retrospective Study

Ezzatollah Rezaei (MD)¹, Arash Beiraghi Toosi (MD)^{1*}, Hassan Aliakbarian (MD)², Hamid Reza Alijani (MD)³, Gholamreza Shariat Gonabadi (MD)³

¹ Associate Professor, Surgical Oncology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

² General Surgeon, Surgical Oncology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

³ General Practitioner, Department of Burns and Reconstructive Surgery, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO	ABSTRACT
<i>Article type:</i> Original article	Background & aim: The incidence of acute burn injuries in pregnant women is very low. Burn injuries during pregnancy are often associated with a high rate of fetal and maternal mortality and morbidity. In this study, we aimed to review the cases of acute burn during pregnancy and evaluate the outcome of these patients in Mashhad, Iran.
<i>Article History:</i> Received: 28-Apr-2015 Accepted: 15-Oct-2015	Methods: This retrospective study was performed using the medical records of 48 pregnant women with thermal injuries over a 13-year period.
<i>Key words:</i> Burns Iran Pregnancy	Results: The results showed that 8 (16.7%), 27 (56.3%), and 13 (27.1%) patients were in the first, second, and third trimesters of pregnancy. Moreover, 14 mothers (29.2%) died, 24 (50.0%) were discharged without any fetal problems, eight (16.7%) had fetal death, 13 (27.1%) had abortion, two (4.2%) had normal vaginal delivery, and one (2.1%) underwent normal caesarean section. Conclusion: The rate of maternal survival in the first and second trimesters was higher than the third one. In the third trimester, pregnancy termination is indicated only after fetal maturation.

► Please cite this paper as:

Rezaei E, Beiraghi Toosi A, Aliakbarian H, Alijani HR, Shariat Gonabadi G. Acute Burns during Pregnancy: A Retrospective Study. Journal of Midwifery and Reproductive Health. 2016; 4(1): 540-543. DOI: 10.22038/jmrh.2016.6123

Introduction

The incidence of acute burn injuries in pregnant women is very low (1); therefore, to delve into this problem, studying large populations over long periods of time is required. Burns during pregnancy are often associated with high rates of fetal and maternal mortality and morbidity. Various factors can affect the rate of mortality and morbidity in burns patients during pregnancy, including depth and size of burn injury, maternal health, age at pregnancy, and simultaneous existence of inhalation and burn injuries (2).

In a study performed by Agrawal, pregnancy did not influence the outcome of acute burn, but it increased the risk of abortion or premature birth (3). In a study by Maghsoudi et al., burn surface area and inhalation injury were the most important predictors of maternal and fetal mortality (4).

Imam Reza Hospital burn center in Mashhad,

Khorasan Razavi province, Iran, serves more than 5.4 million people in an area of 302,000 km² with nearly 1,150 annual admissions (5). In this study, we aimed to review cases of burn injury during pregnancy and evaluate the outcome of these patients in Mashhad, Iran.

Materials and Methods

In this retrospective study, the research protocol was approved by the Ethics Committee of Mashhad University of Medical Sciences. We reviewed the medical records of all cases of burns during pregnancy, who were admitted to Imam Reza Hospital, Mashhad, Iran, during March 2000-March 2013. The obtained data were analyzed regarding age, percentage of total body surface area (TBSA) burned, etiology (self-inflicted or suicidal against incidental burns) and cause of injury (kerosene, hot liquids, or gas explosion),

* Corresponding author: Arash Beiraghi Toosi, Associate Professor, Surgical Oncology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran. Email: beiraghia@mums.ac.ir

gestational age, and fetal and maternal outcome. Patients with mild (first-degree) burn injuries were excluded from the study. For all the included pregnant patients, the data regarding pregnancy laboratory tests, ultrasonography, and gynecologic consultation were recorded.

Results

The results demonstrated that 48 cases of burns during pregnancy were admitted during a

13-year period. On admission, the patients' age ranged from 15 to 43 years, with a mean age of 25.0±6.32 years. In a one-year period (March 2009-March 2010), 439 female burns patients were admitted to our center, 79 (18.0%) of whom died, and 173 (39.4%) were in reproductive age (15-45 years old). The etiology of injury was accidental and self-inflicted in 28 (58.3%) and 20 (41.7%) of the cases, respectively.

Table 1. Age, TBSA^a burns, cause, and maternal and fetal mortality according to the etiology of burns

Etiology	Total No.	Mean age (Y)±SD ¹	TBSA (Mean) ₁	Cause			Maternal mortality ₃	Fetal mortality ² (intrauterine+ abortion)	Abortio n	Fetal death ³
				Kerosene ₃	Gas explosion ₃	Scald				
Accidental	28 (58.3%)	25.1±5.3	31.8%	15 (53.6%)	11 (39.3%)	2 (7.1%)	2 (7.1%)	8 (28.6%)	7 (25.0%)	1 (3.6%)
Suicide	20 (41.7%)	25.8±7.5	63.6%	19 (95.0%)	1 (5.0%)	-	12 (60.0%)	13 (65.0%)	6 (30.0%)	7 (35.0%)
P-value	-	NS ^b	<0.001	0.003	<0.001	-	<0.001	0.012	0.70	0.006

¹ T-student test; ² Chi-square test; ³ Fisher's exact test; ^a TBSA: Total Body Surface Area; ^b NS: non-significant

The injury in 34 (70.8%), 12 (25.0%), and two (4.2%) cases was due to kerosene, gas explosion, and scald. The TBSA ranged between 10% and 95%, with a mean of 45.1±25.49%. The patients' gestational age ranged from 8 to 41 weeks. In addition, 8, 27, and 13 patients were in the first, second, and third trimesters (Table 1).

Twenty-four mothers (50.0%) were discharged without any problems, eight (16.7%) had fetal death, 13 (27.1%) had abortion, two

(4.2%) had normal vaginal delivery, and one (2.1%) had non-complicated cesarean section, totally, 21 (43.8%) fetuses died. Thirty-four patients were eventually discharged, and 14 mothers died. In the other words, in eight cases, both mother and fetus died (intrauterine death). Out of 13 abortions, six mothers died later and seven mothers were discharged. The burn surface area of the dead mothers was more than the survived ones (73.1% vs. 19.0%; P<0.001; Table 2).

Table 2. Maternal and fetal mortality according to TBSA^a burns

TBSA ^a	Mothers ¹		Fetal death ¹ (abortion+intrauterine) (%)
	Total (%)	Died (%)	
<50%	28 (58.3%)	2 (7.1%)	5+1=6 (21.4%)
50-75%	11 (22.9%)	4 (36.4%)	3+4=7 (63.6%)
≥75%	9 (18.8%)	8 (88.9%)	5+3=8 (88.9%)
P-value	-	<0.001	0.001

¹ Chi-square; ^a TBSA: Total Burned Surface Area

The rate of maternal mortality, based on gestational trimester, was as follows: none in the first trimester, 8 out of 27 cases (29.6%) in

the second trimester, and 6 out of 13 cases (46.2%) in the third trimester. These results indicate increased rate of maternal mortality

with increasing gestational age. Table 3 demonstrates the etiology, TBSA, and maternal

and fetal rate of mortality according to gestation trimesters.

Table 3. Etiology, TBSA^a burns, and maternal and fetal mortality according to gestational trimesters

Trimester	Etiology			Mean TBSA (%)	Mother		Fetus			
	Mother's No. (%)	Accidental (%)	Suicidal (%)		Mother's died (%)	Mother's discharge without fetal problems (%)	Normal vaginal delivery (%)	Uncomplicated cesarean section (%)	Abortion / fetal death (%)	Fetal death (%)
1st	8 (16.7%)	6 (75.0%)	2 (25.0%)	35.0%	-	6 (75.0%)	-	-	2 (25%)	-
2nd	27 (56.3%)	15 (55.6%)	12 (44.4%)	51.0%	8 (29.6%)	13 (48.1%)	-	-	10 (37%)	4 (14.8%)
3rd	13 (27.1%)	7 (53.8%)	6 (46.2%)	39.0%	6 (46.2%)	5 (38.5%)	2 (15.4%)	1 (7.7%)	1 (7.7%)	4 (30.8%)
Total	48	24 (58.3%)	20 (41.7%)	45.1%	14 (29.2%)	24 (50%)	2 (4.2%)	1 (2.1%)	13 (27.1%)	8 (16.7%)

^aTBSA: Total Body Surface Area

Discussion

The etiology of burns in 58.3% and 41.7% of the pregnant patients was accidental and suicidal, respectively. Most of the accidental burns took place while cooking. The etiology of burns had a significant relationship with the rate of mortality (3). Among the 28 patients with accidental burn, only two (7.1%) deaths occurred, while one of them had previous history of seizure. Among the 20 patients with self-inflicted burns, there were 12 (60.0%) deaths. The mean TBSA in the accidental and self-inflicted groups was 32.8% and 63.6%, respectively, which were significantly different ($P < 0.001$).

The rate of fetal mortality was higher in the suicidal group (65.0% vs. 28.6%). This result is consistent with those of other similar studies (4, 6). The cause of suicidal burns in the majority of cases was kerosene (95.0%) (4), while different causes of burns (gas explosion and hot liquid) were observed in the accidental group. Maternal rate of mortality had a significant relationship with TBSA (Table 2). Among the 28 patients with less than 50% TBSA burns, only two (7.1%) deaths were observed. This rate in mothers with $\geq 50\%$ and less than 75% TBSA burns was four cases out of 11 (36.4%), and in $\geq 75\%$ TBSA burns the rate was eight out of nine cases (88.9%). One mother (with 85% TBSA burns in the first pregnancy trimester [week 12]) and her fetus survived. These results were much better than those of other studies in which nearly all

mothers with more than 40% (4) or 50% TBSA burns died (7).

The rate of fetal mortality also had a significant relationship with TBSA ($P = 0.001$). In the first group ($< 50\%$ TBSA burns), we had 6 out of 28 (21.4%) cases of fetal death. This rate in the second and third groups was 7 out of 11 (63.6%) and eight out of nine (88.9%) cases, respectively. In this study, it was found that the rate of maternal mortality increased with increasing gestational age. The increased rate of mortality with gestational age is observed when mean TBSA in the third trimester is lower than the second trimester. This pattern proposes that pregnancy and gestational age are risk factors for death in burns patients.

The rate of fetal mortality was the highest in the second trimester, and possibility of cesarean section and vaginal delivery due to fetal viability in the third trimester reduces the rate of fetal mortality at the end of gestational period.

In the first gestational trimester, two out of eight cases (25.0%), in the second trimester 12 out of 27 cases (44.4%), and in the third trimester 6 out of 13 (46.1%) cases were due to suicidal attempts. It suggests that rate of self-inflicted burns increases with increased gestational age. The most common cause of burns was kerosene (73.2%). This finding is in agreement with the results of other similar studies, especially those conducted in Iran (4, 7, 8); although it may be different in other countries (9).

Some studies demonstrating low rate of survival and near complete maternal and fetal mortality in cases of more than 40-50% TBSA burns, recommended early termination of pregnancy in late gestational months (3). In this study, among cases of 50-75% TBSA burns, we had 30% maternal mortality and 60% fetal mortality. Therefore, the increased rate of maternal survival in the first and second trimesters may be attributed to the higher rate of suicidal burns during the third trimester. In the third trimester, pregnancy termination is advised only after fetal maturation.

This study exhibited that suicidal burns comprise a noticeable amount of burns during pregnancy and lead to a high rate of mortality; this result is consistent with those of other studies conducted in Iran (10). This pattern may be different in other countries, even other regions (9, 11-13). In a study done in Iran, the majority (42.7%) of injuries among women of reproductive age were burns followed by lacerations in 32.6% of the cases (14, 15). The limitation of this study was not taking into account the reasons for the attempted suicides due to cultural issues.

In summary, maternal survival in the first and second trimesters was more than the third one. In the third trimester, pregnancy termination was advised only after fetal maturation. Suicidal burns led to higher rates of mortality; additionally, fetal and maternal mortality had a significant relationship with TBSA.

For more precise evaluation of the causes of self-inflicted burn injuries, conducting further studies is required. Investigation of the role of social factors such as addiction, illegitimate pregnancy, and level of education is required. In addition, the role of comorbidities such as diabetes mellitus and inhalation injuries on the rate of mortality should be considered.

Acknowledgements

This study was supported by a grant from the Deputy of Research of Mashhad University of Medical Sciences, Mashhad, Iran.

Conflicts of Interest

The authors declare no conflicts of interest.

References

1. Polko LE, McMahon MJ. Burns in pregnancy. *Obstetrical & Gynecological Survey*. 1998; 53(1): 50-56.
2. Kennedy BB, Baird SM, Troiano NH. Burn injuries and pregnancy. *The Journal of Perinatal & Neonatal Nursing*. 2008; 22(1):21-30.
3. Agarwal P. Thermal injury in pregnancy: predicting maternal and fetal outcome. *Indian Journal of Plastic Surgery*. 2005; 38(2):95-99.
4. Maghsoudi H, Samnia R, Garadaghi A, Kianvar H. Burns in pregnancy. *Burns*. 2006; 32(2):246-250.
5. Rezaei E, Safari H, Motamedolshariati SM, Afzal Aghaei M. Analysis of mortality in a burn center medical. *Journal of Mashhad University of Medical Sciences*. 2009; 52(4):239-243.
6. Karimi H, Momeni M, Momeni M, Rahbar H. Burn injuries during pregnancy in Iran. *International Journal of Gynaecology & Obstetrics*. 2009; 104(2):132-134.
7. Ghotbi SH, Beheshti M. Burn during pregnancy: a five year survey to assess maternal and fetal mortality. *Shiraz E-Medical Journal*. 2006; 7(4):1-5.
8. Maghsoudi H, Pourzand A, Azarmir G. Etiology and outcome of burns in Tabriz, Iran. An analysis of 2963 cases. *Scandinavian Journal of Surgery*. 2005; 94(1):77-81.
9. Haddadin KJ, Haddad SY. Burns in the first trimester of pregnancy. *Annals of Burns and Fire Disasters*. 2003; 16(3):138-139.
10. Mehdizadeh A, Akbarian A, Samareh PP, Tavajjohi S, MacKay RA, Alaghebandan R, et al. Epidemiology of burn injuries during pregnancy in Tehran, Iran. *Annals of Burns and Fire Disasters*. 2002; 15(4):163-169.
11. Othman N, Kendrick D. Epidemiology of burn injuries in the East Mediterranean Region: a systematic review. *BMC Public Health*. 2010; 10(1):83-93.
12. Mabrouk AR, el-Feky AE. Burns during pregnancy: a gloomy outcome. *Burns*. 1997; 23(7):596-600.
13. Akhtar MA, Mulawkar PM, Kulkarni HR. Burns in pregnancy: effect on maternal and fetal outcomes. *Burns*. 1994; 20(4):351-355.
14. Fardiazar Z, Sadeghi-Bazargani H, Mohammadi R. Domestic injuries and suicide among women of reproductive age in Iran. *International Journal of General Medicine*. 2012; 5:547-552.
15. El-Gallal AR, Yousef SM. Burns in pregnancy: a ten-year review of admitted patients. *Annals of Burns and Fire Disasters*. 2002; 15(3):113-115.