

The Relationship between Frequency of Breastfeeding and Sexual Distress in Postpartum Women

Maryam Zamani (MSc)¹, Robab Latifnejad Roudsari (PhD)^{2,3*}, Maryam Moradi (PhD)^{4,3}, Habibollah Esmaily (PhD)^{5,6}

¹ MSc, Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran

² Associate Professor, Nursing and Midwifery Care Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

³ Department of Midwifery, School of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, Iran

⁴ Assistant Professor, Nursing and Midwifery Care Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

⁵ Professor, Social Determinants of Health Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

⁶ Department of Epidemiology and Biostatistics, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article type:
Original article

Article History:
Received: 04-Sep-2017
Accepted: 01-Jan-2018

Key words:
Breastfeeding
Postpartum
Sexual Function

ABSTRACT

Background & aim: Due to frequent waking at night for breastfeeding, nursing mothers are likely to suffer from sleep deprivation, which may lead to sexual dysfunction. To the best of our knowledge, few studies have been conducted on sexual distress in nursing mothers. Thus, we sought to examine the relationship between the frequency of breastfeeding and female sexual distress during postpartum period.

Methods: This descriptive correlational study was conducted on 80 women aged 18-35 years, visiting healthcare centers in Mashhad, Iran, during 2016. To select the participants, multi-stage sampling method was used. The data collection tools included demographic and fertility-related questionnaire, the Depression, Anxiety, and Stress Scale-21 (DASS-21), and the Female Sexual Distress Scale-Revised (FSDS-R). To analyze the data, Kruskal-Wallis H test, Spearman's rank-order correlation, and linear regression model were run in SPSS, version 16.

Results: The mean score of postpartum sexual distress in mothers was 20.43 ± 8.00 . According to the results of the Spearman's rank-order correlation, the frequency of nighttime breastfeeding was directly correlated with maternal sexual distress ($P=0.010$, $r=0.26$). However, there was no correlation between the frequency of daytime breastfeeding sessions and sexual distress ($P=0.976$, $r=0.003$).

Conclusion: Considering the correlation between the frequency of breastfeeding at night and postpartum sexual distress in mothers, it seems necessary to adopt effective counselling approaches to overcome female sexual distress during this period.

► Please cite this paper as:

Zamani M, Latifnejad Roudsari R, Moradi M, Esmaily H. The Relationship between Frequency of Breastfeeding and Sexual Distress in Postpartum Women. Journal of Midwifery and Reproductive Health. 2018; 6(4): 1447-1453.
DOI: 10.22038/jmrh.2018.26061.1285

Introduction

Sexual distress has been defined as depression, anxiety, and stress about sexual activity (1). According to the fifth edition of Diagnostic and Statistical Manual of Mental Disorders (DSM-IV), identification of sexual distress is required to diagnose sexual dysfunctions (2). Moreover, in many women,

sexual function can decrease during pregnancy and the postpartum period (3-6). Sexual problems experienced in the postnatal period can impose great distress on mothers and their partners, which can consequently affect their quality of physical and mental life, as well as their well-being and marital relationship (7).

* *Corresponding author:* Robab Latifnejad Roudsari, Associate Professor, Nursing and Midwifery Care Research Centre, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +989370400607; Email: latifnejadr@mums.ac.ir

In this respect, Shifren et al. (2008) noted that assessing the prevalence of sexual distress to estimate the occurrence of sexual problems requiring clinical interventions was of great importance (8). In addition, postpartum hormonal changes, as well as health recovery and potentially painful and prolonged improvement of the complications of vaginal delivery or cesarean section can influence physical and emotional health status of woman. Due to these factors, couples might struggle to have a healthy, active, and enjoyable sexual life during the postpartum period (9). In this regard, breastfeeding is considered as one of the factors affecting sexual function (10).

The prevalence of sexual dysfunctions and disorders during the postpartum period in women is higher than that in the general population. Moreover, such problems are more common in month 3 and from month 9 to 12 in the postpartum period (10). In this respect, Hicks et al. (2004) estimated the overall prevalence of postnatal sexual dysfunctions in women to be 22-86% (11).

Some investigations have also reported that elevated level of oxytocin due to breastfeeding could have positive effects on sexual function (12, 13). In some cases, high oxytocin levels due to breastfeeding can be associated with low libido (14). Additionally, decreased level of estrogen and high level of prolactin during the postpartum period following breastfeeding can reduce vaginal lubrication and lead to dyspareunia (14). In a report by Schlagintweit et al. (2016), it was found that breastfeeding was not correlated with the frequency and severity of postpartum sexual distress (15). LaMarre et al. (2003) also concluded that breastfeeding mothers compared with bottle-feeders tend to experience more sexual dysfunction, especially loss of libido (7). Moreover, Rowland et al. (2005) showed a strong relationship between breastfeeding and resumption of postpartum sexual intercourse, such that breastfeeding mothers had a later onset of postpartum sexual activity (16). Mothers practicing exclusive breastfeeding can also have more problems with vaginal lubrication and sexual stimulation than others (7). In some studies, sexual function was found

to decrease by 55% in breastfeeding mothers, while it had scaled up by 39% in some other investigations (17). Accordingly, women reporting sexual distress are more likely to suffer from high levels of sexual and marital dissatisfaction (18).

As revealed by the review of the related literature, there is contradictory evidence regarding breastfeeding and sexual function (12, 13, 16, 19, 20, 21). Due to the existing inconsistencies in the studies and given that sexual distress in the postnatal period has not been previously investigated, we aimed to investigate the correlation between the frequency of breastfeeding and maternal sexual distress in the postpartum period.

Materials and Methods

This descriptive correlational study was conducted in 2016 on 80 women aged 18-35 years visiting healthcare centers for neonatal vaccination and care services in Mashhad, Iran. The participants were selected using the multi-stage sampling method during early July to the end of August, 2016. To this end, first, two healthcare centers were randomly selected from five centers located in Mashhad. Then, two affiliate community health centers were randomly selected from each one. Afterwards, convenience sampling method was used to choose the participants from the four community health centers of Imam Hassan (AS), Ghandkooh, 14 Maasoum, and Imam Reza (AS). The sample size was calculated with 95% confidence interval and 80% power.

The inclusion criteria were women having Iranian nationality, residing in the city of Mashhad, being aged 18-35 years, having high school diploma or higher education, being married, being their spouse's only partner, living with the spouse, being in the 3 to 12 month postpartum, having a healthy and full-term singleton pregnancy, having routine sexual relationships, reporting at least one untreated sexual disorder within the last two months, and obtaining the Depression, Anxiety, and Stress Scale-21 (DASS-21) scores less than 13, 9, and 17 in the domains of depression, anxiety, and stress (normal range), respectively.

The exclusion criteria included suffering from drug or alcohol addiction, taking medications affecting sexual function, experiencing a stressful incident over the past six months, suffering from specific medical conditions, having a history of pelvic surgery or radiation to their genitalia, suffering from certain mental and psychological problems, and experiencing postpartum complications in their recent delivery.

After receiving permission from the Ethics Committee of Mashhad University of Medical Sciences and to conduct sampling and data collection, the researcher offered a letter of introduction from the School of Nursing and Midwifery to the respective authorities in the healthcare centers and then made the necessary arrangements. After introduction and explaining the study objectives for the participants, they were included in the study if they met the inclusion criteria. The respondents were also ensured of the confidentiality of their data, and then signed the informed consent form. As soon as the participants received healthcare services from the given centers, they were guided to a quiet room in the centers to observe their privacy. After the necessary information was provided on how to respond to the questionnaires, the demographic and fertility-related questionnaire, the DASS-21, and the FSDS-R were completed.

The demographic and fertility-related questionnaire contained five sections including personal, marital, pregnancy and delivery, postpartum and sexual data. The validity of this instrument was determined through content validity

In this study, the short form of the DASS-21 comprised of 21 items using a 4-point Likert type scale from zero to three was used. The highest score in each of the mentioned sub-groups was 21. Obtaining scores 0-14, 15-18, and 19-21 indicated normal, mild, and moderate and severe stress, respectively. Moreover, earning a score of 0-7 represented normal anxiety, scores of 8-9 showed mild anxiety, and scores of 10-21 indicated moderate or severe anxiety. Furthermore, scores of 0-9 showed normal depression, scores of 10-13 indicated mild depression, and scores of 14-21

represented moderate or severe depression. The validity and reliability of the DASS-21 was confirmed by Sahebi et al. in Iran in 2002 (22). The reliability of this tool was also determined in this study using the Cronbach's alpha method ($\alpha=0.91$).

The Female Sexual Distress Scale-Revised (FSDS-R) comprised of 13 items examining various dimensions of female sexual activity. All the items were scored based on a five-point Likert type scale from 0 (never) to 4 (always). The overall score was obtained by aggregating all the item scores. Moreover, scores greater than and equal to 11 represented sexual distress. The validity and reliability of this tool was measured by Derogatis et al. (2002) (23). In Iran its validity and the reliability was also determined by Ghassami et al. (2014) (24). It should be noted that the reliability of this tool was confirmed in this study via Cronbach's alpha method ($\alpha=0.86$).

To determine the relationship between the frequency of breastfeeding sessions and maternal sexual distress, Spearman's rank-order correlation and linear regression model were employed. The normality test was also run using the Kolmogorov-Smirnov test in SPSS, version 16. *P*-value less than 0.05 was considered statistically significant.

Results

The mean age of the women and their spouses were 29.5 ± 4.2 and 34.3 ± 5.7 years, respectively. Overall, 52% ($n=42$) of these mothers had high school diploma, 46.3% ($n=35$) of them had a history of vaginal delivery, and 53.7% ($n=45$) had cesarean section. Menstruation had also returned in 58.8% ($n=46$) of the mothers, and 55% ($n=43$) of them were practicing exclusive breastfeeding (Table 1). The average days of menstruation in the postpartum period were 58.8 ± 48.7 . The mean number of gravidity was 1.7 ± 0.7 , the mean number of deliveries was 1.6 ± 0.6 , and the mean number of children was 1.6 ± 0.5 . Moreover, the mean score of sexual distress was 20.43 ± 8.00 . The mean scores of the depression, anxiety, and stress were 11.06 ± 3.28 , 14.87 ± 2.83 , and 14.53 ± 3.68 , respectively.

According to the results of the Kruskal-Wallis H test, sexual distress was correlated with the

type of infant's nutrition ($P=0.001$).

Table 1. Demographic and fertility characteristics of women in the postpartum period and their relationship with sexual distress

Variable	Frequency (%)
Education	
High school diploma	42 (52.5)
Associate degree	9 (11.3)
Bachelor degree	29 (36.3)
* $P=0.253$	
Frequency of daytime breastfeeding	
1-2 times	6 (7.5)
3-4 times	8 (10.0)
4-5 times	16 (20.0)
More than 6 times	50 (62.5)
* $P=0.053$	
Frequency of nighttime breastfeeding	
1-2 times	25 (31.3)
3-4 times	41 (51.3)
4-5 times	9 (11.3)
More than 6 times	5 (6.3)
* $P=0.051$	
Income	
Not given	4 (5.0)
Enough	59 (75.0)
Less than enough	17 (20.0)
* $P=0.937$	
Route of delivery	
Vaginal	35 (46.3)
Cesarean-section	45 (53.7)
* $p=0.773$	
Menstruation return	
Yes	46 (58.8)
No	34 (42.2)
Contraceptive method	
Linstrol	7 (8.8)
Intrauterine devices (IUD)	4 (6.3)
Condom	28 (35.0)
Interrupted sex	12 (13.8)
Vasectomy	8 (10.0)
No sex	21 (26.3)
* $p=0.246$	
Infant's nutrition	
Exclusive breastfeeding	43 (55.0)
Bottle-feeding	5 (6.3)
Exclusive breastfeeding + bottle-feeding	6 (1.3)
Supplementary foods + breastfeeding	31 (37.5)
* $P=0.001$	

*Kruskal-Wallis H test

According to this test, it was found that sexual distress had a borderline relationship with the frequency of nighttime breastfeeding ($P=0.051$)

and those during the day ($P=0.053$; Table 1). According to the findings of the Kolmogorov-Smirnov test, sexual distress scores had an

abnormal distribution; thus, Spearman's rank-order correlation was used. The results of the Spearman's rank-order correlation similarly showed no direct correlation between the frequency of daytime nursing and sexual distress ($P=0.976$, $r=0.003$). However, according to the findings of the Spearman's rank-order correlation, there was a direct correlation between the frequency of breastfeeding at night and female sexual distress, such that the frequency of nursing at night increased sexual distress in mothers ($P=0.010$, $r=0.286$). Also,

Spearman's rank-order correlation revealed a direct correlation between sexual distress and depression, that is, the higher sexual distress, the more severe the level of depression ($P=0.008$, $r=0.294$). Considering the results of the Spearman's rank-order correlation, no statistically significant relationship was observed between the frequency of nursing frequency during the day as well as breastfeeding times at night and the levels of depression ($P=0.676$), anxiety ($P=0.851$), and stress ($P=0.583$; Table 2).

Table 2. Spearman's rank-order correlation of sexual distress and frequency of daytime and nighttime breastfeeding and depression, anxiety, stress

Sexual distress	Frequency of daytime breastfeeding	Frequency of nighttime breastfeeding	Depression	Anxiety	Stress
Correlation coefficient	0.003	0.286	0.294	0.172	0.085
P-value	0.976	0.010	0.008	0.127	0.454

Given the results of the linear regression model, there was a linear relationship between the frequency of breastfeeding at night and maternal sexual distress ($P=0.001$), such that increased frequency of nursing at night led to a rise by 0.393 in the mean score of sexual

distress. However, according to the linear regression model, there was no significant association between the frequency of daytime breastfeeding and female sexual distress ($P=0.167$; Table 3).

Table 3. Linear regression model for the variables of frequency of daytime and nighttime breastfeeding

Predictive variables	Regression coefficient	Standard deviation	Standardized regression coefficient	Test statistic	Pa-value
Frequency of nighttime breastfeeding	3.818	1.134	0.393	3.365	0.001
Frequency of daytime breastfeeding	-1.387	0.987	-0.163	-1.397	0.167

Discussion

The present study was conducted to determine the link between the frequency of breastfeeding and maternal sexual distress during the postpartum period. The findings of this study showed a direct correlation between the frequency of nighttime nursing frequency and postpartum sexual distress in mothers. In this regard, increased frequency of breastfeeding at night was often accompanied by higher levels of sexual distress. However, there was no direct relationship between the frequency of breastfeeding during the day and maternal sexual distress.

It should be noted that studies on sexual distress, particularly, on women's sexual

distress during the postpartum period, are few. In this respect, Lagaert et al. (2017) reported that breastfeeding mothers could suffer from sleep deprivation due to the frequency of nursing at night, which could be accompanied by sexual dysfunction (25); their findings are congruent with our results. It seems that maternal fatigue can be considered one of the most important factors affecting mothers' sexual performance. The results of a study by Shirvani et al. (2011) also demonstrated a direct correlation between maternal fatigue and sexual dysfunction (26). In addition, Rowland et al. (2005) reported that the level of prolactin in mothers could go up after breastfeeding, which could decrease gonadotropin level, and consequently, lead to reduced level of estrogen.

Vaginal dryness could also occur during the postpartum period because of estrogen reduction followed by dyspareunia, that is, difficult and painful sexual intercourse (16).

Likewise, the results of the investigation by Khosravi Anbaran et al. (2014) suggested that postpartum sexual function in women is correlated with breastfeeding, so those who exclusively breastfeed their babies could suffer from more sexual problems (27).

In a review titled as "Breastfeeding and postpartum maternal sexual functioning: A review", LaMarre et al. (2003) found that breastfeeding mothers reported a decrease in their sexual function, especially libido, compared with those who bottle-feed their babies (7). According to Abdool et al. (2009) most studies found that breastfeeding had a significant relationship with libido (28). According to the results of this study, there was no statistically significant relationship between the frequency of breastfeeding at night and those during the day and depression, anxiety as well as stress. Rowland et al. (2005) did not find any evidence of mood changes caused by the negative effects of breastfeeding on sexual function (16). Zamani et al. citing Huang et al. (2006) discussed that variability of the relationship between sexual function and postpartum depression is a culture-dependent issue, which may vary from country to country (29). They also suggested that breastfeeding mothers could benefit from talking to their healthcare providers about sexual and breastfeeding issues. Thus, healthcare providers should reassure mothers that changes in their sexual activity and levels of sexual satisfaction during the postpartum period are expected (16).

Among the limitations of this study was the effect of personal motivations of the participants to respond to the questionnaires due to the self-reporting nature of the given tools, as well as the limited sample size. We suggest investigating the impact of breastfeeding on postpartum sexual distress in future studies using larger sample size.

Conclusion

Considering the relationship between the frequency of nighttime breastfeeding and postpartum sexual distress in mothers, it seems

necessary to employ effective counselling approaches to moderate female sexual distress in this period.

Acknowledgements

This research study was part of a larger clinical trial approved by the Ethics Committee of Mashhad University of Medical Sciences and registered at the Iranian Registry of Clinical Trials with the code number of IRCTID: IRCT2016062228575N. We wish to thank the Deputy of Research of Mashhad University of Medical Sciences for their financial support (project code: 941778). We also sincerely appreciate all the mothers for their in this study.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Stephenson KR, Meston CM. Differentiating components of sexual well-being in women: Are sexual satisfaction and sexual distress independent constructs? *The Journal of Sexual Medicine*. 2010; 7(7):2458-2468.
2. American Psychiatric Association. Task force on DSM-IV. *DSM-IV options book: work in progress* (7.1.91). New York: American Psychiatric Press; 1991.
3. Safarinejad MR. Female sexual dysfunction in a population-based study in Iran: prevalence and associated risk factors. *International Journal of Impotence Research*. 2006; 18(4):382-395.
4. West SL, D'Aloisio A, Agans RP, Kalsbeek WD, Borisov NN, Thorp JM. Prevalence of low sexual desire and hypoactive sexual desire disorder in a nationally representative sample of US women. *Archives of Internal Medicine*. 2008; 168(13):1441-1449.
5. Warnock JK, Clayton A, Croft H, Segraves R, Biggs FC. Women's sexual health: comparison of androgens in women with hypoactive sexual desire disorder: those on combined oral contraceptives (COCs) vs. those not on COCs. *The Journal of Sexual Medicine*. 2006; 3(5):878-882.
6. Dennerstein L, Guthrie JR, Hayes RD, DeRogatis LR, Leher P. Sexual function, dysfunction, and sexual distress in a prospective, population-based sample of mid-aged, Australian-born women. *The Journal of Sexual Medicine*. 2008; 5(10):2291-2299.
7. LaMarre AK, Paterson LQ, Gorzalka BB. Breastfeeding and postpartum maternal sexual functioning: A review. *The Canadian Journal of Human Sexuality*. 2003; 12(3/4):151.

8. Shifren JL, Monz BU, Russo PA, Segreti A, Johannes CB. Sexual problems and distress in United States women: Prevalence and correlates. *Obstetrics & Gynecology*. 2008; 112(5):970-978.
9. Alves M, Vieira R. T10-O-01 facts that influence the feminine sexuality after the childbirth. *Sexologies*. 2008; 17:S139.
10. Tork Zahrani S, Banaei M, OZgoli G, Azad M. Investigation of the postpartum female sexual dysfunction in breastfeeding women referring to healthcare centers of Bandar Abbas. *The Iranian Journal of Obstetrics, Gynecology, and Infertility*. 2016; 19(35):1-12. (Persian)
11. Hicks TL, Goodall SF, Quattrone EM, Lydon-Rochelle MT. Postpartum sexual functioning and method of delivery: Summary of the evidence. *Journal of Midwifery & Women's Health*. 2004; 49(5):430-436.
12. Avery MD, Duckett L, Frantzich CR. The experience of sexuality during breastfeeding among primiparous women. *Journal of Midwifery & Women's Health*. 2000; 45(3):227-237.
13. McBride HL, Kwee JL. Sex after baby: women's sexual function in the postpartum period. *Current Sexual Health Reports*. 2017; 9(3):142-149.
14. McBride HL, Olson S, Kwee J, Klein C, Smith K. Women's postpartum sexual health program: a collaborative and integrated approach to restoring sexual health in the postpartum period. *Journal of Sex & Marital Therapy*. 2017; 43(2):147-158.
15. Schlagintweit HE, Bailey K, Rosen NO. A new baby in the bedroom: frequency and severity of postpartum sexual concerns and their associations with relationship satisfaction in new parent couples. *Journal of Sexual Medicine*. 2016; 13(10):1455-1465.
16. Rowland M, Foxcroft L, Hopman WM, Patel R. Breastfeeding and sexuality immediately post partum. *Canadian Family Physician*. 2005; 51(10):1366-1367.
17. Rezaei N, Azadi A, Sayehmiri K, Valizadeh R. Postpartum sexual functioning and its predicting factors among Iranian women. *The Malaysian Journal of Medical Sciences*. 2017; 24(1):94.
18. Vannier SA, Rosen NO. Sexual distress and sexual problems during pregnancy: Associations with sexual and relationship satisfaction. *The Journal of Sexual Medicine*. 2017; 14(3):387-395.
19. Faisal-Cury A, Menezes PR, Quayle J, Matijasevich A, Diniz SG. The relationship between mode of delivery and sexual health outcomes after childbirth. *The Journal of Sexual Medicine*. 2015; 12(5):1212-1220.
20. Barbara G, Pifarotti P, Facchin F, Cortinovis I, Dridi D, Ronchetti C, et al. Impact of mode of delivery on female postpartum sexual functioning: Spontaneous vaginal delivery and operative vaginal delivery vs. Cesarean-section. *The Journal of Sexual Medicine*. 2016; 13(3):393-401.
21. Dahlen HG, Barclay LM, Homer CS. Processing the first birth: journeying into 'motherland'. *Journal of Clinical Nursing*. 2010; 19(13-14):1977-1985.
22. Sahebi A, Asghari MJ, Salari RS. Validation of depression anxiety and stress scale (DASS-21) for an Iranian population. *Iranian Psychologists*. 2005; 4(1):299-313.
23. Derogatis LR, Rosen R, Leiblum S, Burnett A, Heiman J. The female sexual distress scale (FSDS): initial validation of a standardized scale for assessment of sexually related personal distress in women. *Journal of Sex & Marital Therapy*. 2002; 28(4):317-330.
24. Ghassami M, Asghari A, Shaeeri MR, Soltaninejad Z, Safarinejad MR. Psychometric properties of the Female Sexual Distress Scale-Revised among a sample of non-clinical Iranian women. *International Journal of Psychiatry in Clinical Practice*. 2014; 18(4):293-299.
25. Lagaert L, Weyers S, Van Kerrebroeck H, Elaut E. Postpartum dyspareunia and sexual functioning: a prospective cohort study. *The European Journal of Contraception & Reproductive Health Care*. 2017; 22(3):200-206.
26. Ahmad Shirvani M, Bagheri Nesami M. Sexual dysfunction and related factors among breast feeding women. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. 2011; 14(5):38-44. (Persian)
27. Khosravi Anbaran Z, Baghdari N, Pourshirazi M, Karimi A. The relationship between women's postpartum sexual function and infant feeding. *Iranian Journal of Neonatology*. 2014; 5:2-3.
28. Abdool Z, Thakar R, Sultan AH. Postpartum female sexual function. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*. 2009; 145(2):133-137.
29. Zamani M., Latifnejad Roudsari R, Moradi M, Esmaily H. Effect of Sexual Counseling on Stress, Anxiety, and Depression in Women during Postpartum Period. *Evidence Based Care*. 2017, 7(2):17-26.