

# Prevalence of Primary Ovarian Insufficiency, Early Menopause and their Associated Factors among Menopausal Women: Results from the AZAR Cohort study in Iran

Sepideh Mashayekh-Amiri (PhD)<sup>1,2</sup>, Elnaz Faramarzi (PhD)<sup>3</sup>, Mojgan Mirghafourvand (PhD)<sup>4,5\*</sup>

<sup>1</sup> Assistant Professor, Counseling and Reproductive Health Research Centre, Golestan University of Medical Sciences, Gorgan, Iran

<sup>2</sup> Department of Midwifery, Faculty of Nursing and Midwifery, Golestan University of Medical Sciences, Gorgan, Iran

<sup>3</sup> Associate Professor, Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>4</sup> Professor, Social Determinants of Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>5</sup> Menopause Andropause Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

## ARTICLE INFO

*Article Type:*  
Original article

*Article History:*  
Received: 04-Apr-2024  
Accepted: 08-Sep-2025

*Key words:*  
Primary Ovarian Failure  
Early Menopause  
Age at Menopause  
AZAR Cohort Study

## ABSTRACT

**Background & aim:** Primary ovarian insufficiency (POI) and early menopause are among the primary causes of female infertility. Thus, the present study evaluated the prevalence of POI and early menopause and identifies the associated factors in postmenopausal women in Shabestar, Iran.

**Methods:** This cross-sectional study was carried out using data from the AZAR cohort study (ACS), which was conducted in Shabestar, Iran, from October 2014 to January 2017. Among 15,006 participants, fundamental data for 8,290 women were extracted, of whom 3,491 were in menopausal period. Data were collected using a structured and pre-designed questionnaire. To identify the prevalence of POI and associated factors, univariable and multivariable logistic regression were used.

**Results:** This study found that the prevalence of POI, early menopause, and age at natural menopause was 6.1%, 27.6%, and 66.3%, respectively. Multivariable logistic regression results, after adjusting for all potential confounding factors, revealed that the odds of POI were significantly lower in women with parity  $\geq 3$ , overweight women, and women with late menarche. The odds of POI were also considerably higher in smokers. The results also revealed that underweight women had more than 4 times higher odds of early menopause.

**Conclusion:** The results revealed that parity, age at menarche, overweight, and smoking had a statistically significant relationship with POI. Due to the increasing prevalence of POI and early menopause, screening to identify the associated factors by healthcare providers will help to provide early interventions.

### ► Please cite this Paper as:

Mashayekh Amiri S, Faramarzi E, Mirghafourvand M. Prevalence of Primary Ovarian Insufficiency, Early Menopause and their Associated Factors among Menopausal Women: Results from the AZAR Cohort study in Iran. Journal of Midwifery and Reproductive Health. 2026; 14(3): 5519-5531. DOI: 10.22038/jmrh.2026.82319.2468

## Introduction

Menopause, as the end point of women's reproductive period, is considered a natural transitional process in women's lives (1). Permanent cessation of menstrual cycles for 12 consecutive months without any apparent psychological or pathological reason (such as bilateral oophorectomy, chemotherapy, or radiotherapy) is called natural menopause (2). The mean age of natural menopause is

estimated to be 51.4 years, but it varies based on the geographical region (3).

Menopausal age is considered a very significant biomarker and indicator of ovarian function. Menopause before the age of 40 is classified as primary ovarian insufficiency (POI). Menopause between the ages of 40 and 45 is called early menopause (4). In the United States, about 5 percent of women experience menopause between the ages of 40 and 45 years. Also, 1% of women experience

\* Corresponding author: Mojgan Mirghafourvand, Professor, Social Determinants of Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. Tel; 0098914306121; Email: mirghafourvand@gmail.com



menopause before the age of 40 (5). The results of a meta-analysis study revealed that about a quarter of Iranian women experience menopause earlier than the age of 45 (6). The results of a survey from 9 cohort studies on 51,450 postmenopausal women reported the prevalence of POI and early menopause as 0.2% and 7.6%, respectively (7). The results of a systematic review study in Iran revealed that the prevalence of POI and early menopause is 3.7% and 12.2%, respectively (6).

Primary ovarian insufficiency is defined as the cessation of menstruation with a lack of serum estrogen and an increase in follicle-stimulating hormone (FSH) more than mIU/L40, at the age of less than 40 years, with approximately 2 standard deviations lower than the mean age of natural menopause (8). Therefore, POI occurs at least 10 years before the mean menopause age of 51.4 years (9). The manifestations of POI include one of the symptoms of secondary amenorrhea or oligomenorrhea, infertility, and estrogen deficiency. POI and early menopause can occur spontaneously and be induced (10).

The causes of POI are very heterogeneous. The guideline development group of the European Society of Human Reproduction and Embryology (ESHRE) has classified the causes of POI into 6 categories: chromosomal or genetic defects, iatrogenic causes, infections, autoimmune disorders, environmental factors, and idiopathic factors (11). The results of a study of more than 11,000 Australian women revealed that women with POI were almost three times more likely to develop multiple diseases in their sixties than women with natural menopause (12). In other words, women with POI experience the effects of estrogen deprivation for a longer period, which is associated with many complications.

POI and early menopause are associated with many psychosocial and physiological problems, including increased risk of osteoporosis (13), cardiovascular disease (14), type 2 diabetes (15), Alzheimer's (16), early death (17), depression (18), and low quality of life (19). Several factors are involved in determining the time of menopause and its deviation from the normal range. The results of a study by Vatankhah et al. (2023) on factors related to

menopause showed that age, height, smoking, history of diabetes, hypertension, thyroid disease, depression, and use of hormonal contraceptives had a significant relationship with menopause (20).

The results of the Tehran lipid and glucose study (TLGS) using the Accelerated Failure Time (AFT) model showed that, compared with time of menopause in normal-weight women, it decreased by 0.09 and 0.03 years in underweight and overweight women, respectively. Also, the natural menopause time increased by 0.12 years in women who used oral contraceptives for more than 6 months (21). Since women in developing countries experience menopause earlier than women in developed countries (22), and since reported risk factors for early menopause, including female smoking, obesity, and early menarche, are increasing, it is crucial to determine the prevalence of POI and early menopause and identify the factors associated with it.

Although a significant relationship between POI and early menopause and the health status and quality of life of women has been reported, there is currently a dearth of knowledge about the prevalence of POI and early menopause and its associated factors, especially in the Asian population. Thus, identifying the aspects related to POI and early menopause, especially those that are modifiable, may be crucial for planning health and prevention interventions and improving the quality of life of postmenopausal women. Hence, this study evaluated the prevalence of POI and early menopause and identified the associated factors in postmenopausal women in Shabestar, Iran.

## Materials and Methods

This cross-sectional study was conducted using data from the AZAR cohort study (ACS), as part of the prospective epidemiological research studies in Iran (PERSIAN cohort in Shabestar, one of the cities in East Azerbaijan province, Iran (23, 24). ACS is a population-based prospective cohort that aims to investigate non-communicable diseases (NCDs), including cardiovascular, gastrointestinal, liver, kidney, metabolic, respiratory disorders, and cancers. It was initiated in October 2014 and ended in January 2017. Details of this cohort study has been published elsewhere (25).

The inclusion criteria of the study included living in Shabestar city for at least 9 months and being aged 35 to 70 years. People with severe mental and physical diseases were excluded from the study. For sampling, almost all eligible people in small towns and villages were invited to participate in the study. Also, in bigger cities, more than 60% of the target population was asked (25). First, the list and medical records of all people aged 35 to 70 years were prepared by Shabestar Health Centers (East Azerbaijan Province, Iran).

After checking the inclusion and exclusion criteria, if the women were willing to participate in the study, written informed consent was obtained from them. Among 15006 participants, the basic data of 8290 women were extracted. In this step, women who never experienced menstruation in their lifetime ( $n=4$ ) were excluded from the study. Finally, 8286 women entered the present study. Among them, 3491 were in menopausal period and provided reliable information about their menopause. According to the World Health Organization (WHO) classification, menopause is defined as the absence of spontaneous menstrual bleeding for more than 12 months without any other pathological or physiological cause (26). The time after the 12 months of no menstrual bleeding was considered the age of natural menopause (ANM). POI was defined as menopause before the age of 40, and early menopause was defined as menopause between the ages of 40 and 45 years (4).

Data were collected by trained interviewers using a valid and reliable questionnaire administered face-to-face (25). It included information related to socio-demographic and anthropometric characteristics such as age, age at menarche, BMI, gravidity, number of abortions, marital status, and educational level, quintiles of wealth index, smoking status, oral contraceptives, fertility drugs, infertility history, tubectomy, and hysterectomy history. To determine anthropometric indicators, BMI was calculated as weight (kg) divided by height (m) squared after measuring each participant's height and weight. BMI classification was used based on the World Health Organization. Accordingly, participants were classified into four different categories: (1) underweight (BMI

< 18.5 kg/m<sup>2</sup>), (2) normal weight (BMI: 18.5-24.9 kg/m<sup>2</sup>), (3) overweight (BMI: 25.0-29.9 kg/m<sup>2</sup>, and (4) obese (BMI  $\geq$  30 kg/m<sup>2</sup>). Regarding smoking, smokers were those who continuously smoke at least one cigarette per day for more than half a year. Ex-smokers were people who used to smoke in the past but do not smoke now. Non-smokers had no history of smoking, and other tobacco users were those who smoked hookah and chewed snuff.

Socioeconomic status was estimated based on Wealth Score Index (WSI) using Multiple Correspondence Analysis (MCA) according to the variables such as access to a dishwasher, access to a computer, access to a car, price of a car, owning a mobile phone, number of international trips in a lifetime, and household amenities (e.g., number of rooms, ownership type). The participants of this study were divided into five classes based on WSI, ranging from poorest to richest. Fertility history information included age at menarche, gravidity, parity, number of abortions, use of oral contraceptive pills (OCP), history of taking fertility drugs, history of taking drugs for infertility, hysterectomy, and tubectomy.

To analyse data, SPSS statistical software (Version 25.0) (IBM Corp, Armonk, NY, USA) was used. Socio-demographic characteristics and the prevalence of POI and early menopause among participants were described using descriptive statistics, including frequency, percentage, mean, and standard deviation (SD). To estimate the prevalence of POI and early menopause, the probability of menopause events was used based on age. Women's menopausal age was classified into 3 groups with 5-year intervals based on available sources. Then, the menopause rate in each age group was estimated. Skewness and Kurtosis were used to determine the normality of the quantitative data. They indicated that the data followed a normal distribution. To determine the relationships between POI and early menopause and various variables, bivariate tests, including univariate logistic regression, were used to estimate crude odds ratios (cORs). Then, variables with p-values < 0.2 were entered into the multivariable logistic regression model to control for confounding. In the statistical analysis, the age at natural menopause was used

as the reference category. The results of multivariable logistic regression were presented as adjusted odds ratios (aORs) with 95% confidence intervals (CIs). P-value <0.05 was considered statistically significant.

## Results

In the present study, as part of the ACS study, 8286 participants were enrolled from Oct 2014 to Jan 2017. Among them, 3491 were menopausal. The mean age (SD) of the participants and their mean age of menopause (SD) in this study were 49.2 (9.3) and 47.4 (5.2), respectively. Almost over half of the women (60%) were over 45 years old. Regarding marital status, the majority of women (87.8%) were married. More than half of women (59.3%) menstruated for the first time in the normal age range (12-14 years). The majority of them (84.5%) were overweight or obese. More than two-thirds of the participants (71.5%) reported parity  $\geq 3$ . The majority of them (94%) had less than a diploma. Also, more than two-thirds of them (73.8%) had a history of using oral contraceptive methods. Based on the WSI index, more than a quarter of women (27.5%) had a poor socioeconomic status. Also, the majority of women (92.3%) did not report a history of using drugs for infertility treatment. Table 1 presents other details related to demographic characteristics and anthropometric measures.

**Table 1.** Socio-demographic characteristics among studied women (N=8286)

Characteristic	N (%)
<b>Age (Years)</b>	
<40	1472 (17.8)
40-45	1841 (22.2)
>45	4973 (60.0)
<b>Age at menarche (Years)</b>	
Early<12	648 (7.8)
Normal (12-14)	4911 (59.31)
Late>14	2727 (32.9)
<b>BMI (kg/m<sup>2</sup>)</b>	
Underweight (<18.5)	(0.4) 33
Normal (18.5-24.9)	(15.1) 1249
Overweight/Obese (>25)	(84.5) 6996
<b>Gravidity</b>	
0	(5.4) 447
1-2	(23.1) 1918
$\geq 3$	(71.5) 5921

Characteristic	N (%)
<b>Number of abortions</b>	
No abortion	(62.6) 4096
One	(24.4) 1912
$\geq 2$	(13.0) 1021
<b>Marital status</b>	
Single	(2.4) 199
Married	(87.8) 7271
Widow/divorced	(9.8) 816
<b>Educational level</b>	
Illiterate	(23.2) 1921
Primary school	(40.9) 3393
Intermediate/ High school	(29.9) 2477
University	(6.0) 495
<b>Quintiles of wealth index</b>	
1 (poorest)	(27.5) 2280
2	(17.8) 1474
3	(19.7) 1629
4	(20.6) 1703
5 (richest)	(14.5) 1200
<b>Smoking status</b>	
Never	(98.8) 8188
Ex-smoker	(0.5) 38
Smoker	(0.5) 39
Smoker of other tobacco products	(0.2) 21
<b>Oral contraceptives</b>	
Yes	(73.8) 6119
No	(26.2) 2167
<b>Fertility drugs</b>	
Yes	(5.6) 451
No	(94.4) 7636
<b>Infertility history</b>	
Yes	(7.7) 626
No	(92.3) 7461
<b>Tubectomy</b>	
Yes	(24.6) 2036
No	(75.4) 6250
<b>Hysterectomy</b>	
Yes	(4.2) 352
No	(95.8) 7934

BMI: body mass index

## The prevalence of POI and early menopause

The results of this study revealed that the prevalence of POI (age less than 40 years), early menopause (age between 40 and 45), and menopause at natural age (age more than 45 years) was 6.1%, 27.6%, and 66.3%, respectively (Table 2).

**Table 2.** The prevalence of POI and early menopause (N=3491)

Menopause status	The prevalence of POI and early menopause, N (%)
POI (<40)	213 (6.1)
Early (40-45)	964 (27.6)
Natural (>45)	2314 (66.3)

POI: Primary ovarian insufficiency

**Factors associated with POI and early menopause based on bivariate and multivariable analyses**

Based on the univariate logistic regression test results, a statistically significant relationship was found between the variables of BMI (P=0.020), gravidity (P<0.001), parity (P<0.001), smoking (P<0.001), use of fertility drugs (P=0.036), history of infertility (P=0.005), and POI. Also, a statistically significant relationship was observed among BMI (P=0.031), gravidity (P= 0.004), parity (P= 0.012),

use of fertility drugs (P= 0.020), history of infertility (P= 0.041), and early menopause (Table 3).

Then, the variables that had P<0.2 were entered into the multivariable logistic regression model. After adjusting the demographic characteristics of the participants, the results of multivariable logistic regression showed that the odds of POI was significantly lower in women with parity≥3 (aOR 0.37; 95% CI 0.19 to 0.69, P=0.002), overweight women (aOR 0.61; 95% CI 0.39 to 0.96, P= 0.033), and women with late menarche (aOR 0.67; 95% CI 0.49 to 0.92, P= 0.014). In other words, women with parity≥3, overweight women, and women with a history of late menarche had lower odds of having POI by 63%, 39%, and 33%, respectively. Also, the odds of POI were significantly higher in smoking women (aOR 7.85; 95% CI 2.60 to 23.67, P<0.001). In other words, smoking increased the odds of POI almost 8 times (Table 4).

**Table 3.** Univariable logistic regression analysis for factors associated with POI and EM (N=1177)

Variable	POI (<40) (N=213)		EM (40-45) (N=964)	
	cOR (95% CI)**	P-Value	cOR (95% CI)**	P-Value
<b>Age of menarche (Years)</b>				
Normal (12-14) (Ref)	1		1	
Early <12	0.97 (0.58-1.63)	0.921	1.16 (0.88-1.544)	0.270
Late >14	0.74 (0.55-1.01)	0.060	1.07 (0.91-1.25)	0.380
<b>BMI (kg/m<sup>2</sup>)</b>				
Normal (18.5-24.9) (Ref)	1		1	
Underweight (<18.5)	2.84 (0.28-28.12)	0.370	4.67 (1.15-18.97)	0.031
Overweight (25-29.9)	0.60 (0.39-0.94)	0.020	0.95 (0.75-1.21)	0.708
Obese (≥30)	0.85 (0.57-1.27)	0.450	0.94 (0.74-1.19)	0.627
<b>Gravidity</b>				
<3 (Ref)	1		1	
≥3	0.53 (0.36-0.77)	<0.001	0.71 (0.57-0.89)	0.004
<b>Parity</b>				
<3 (Ref)	1		1	
≥3	0.49 (0.35-0.68)	<0.001	0.77 (0.63-0.94)	0.012
<b>Number of abortions</b>				
No abortion (Ref)	1		1	
One	1.02 (0.73-1.41)	0.886	0.94 (0.78-1.12)	0.511
≥2	0.74 (0.47-1.16)	0.196	0.97 (0.78-1.20)	0.780
<b>Marital status</b>				
Single (Ref)	1		1	
Married	1.64 (0.21-12.40)	0.628	0.73 (0.33-1.59)	0.430
Widow/divorced	1.73 (0.22-13.39)	0.595	0.81 (0.37-1.80)	0.619
<b>Educational level</b>				

Variable	POI (<40) (N=213)		EM (40-45) (N=964)	
	cOR (95% CI)**	P-Value	cOR (95% CI)**	P-Value
Illiterate	2.71 (0.65-11.26)	0.170	1.06 (0.65-1.73)	0.796
Primary school	2.98 (0.71-12.44)	0.133	1.06 (0.65-1.74)	0.801
Intermediate/ High school	2.42 (0.56-10.39)	0.232	0.82 (0.49-1.38)	0.471
University (Ref)	1		1	
<b>Quintiles of wealth index</b>				
1 (poorest)	1.378 (0.83-2.26)	0.207	1.24 (0.95-1.60)	0.102
2	0.98 (0.56-1.69)	0.947	1.05 (0.80-1.39)	0.683
3	1.26 (0.73-2.18)	0.397	1.02 (0.76-1.37)	0.861
4	1.0 (0.57-1.92)	0.866	0.91 (0.66-1.25)	0.563
5 (richest) (Ref)	1		1	
<b>Smoking status</b>				
Never (Ref)	1		1	
Ex-smoker	0.97 (0.22-4.14)	0.967	0.51 (0.19-1.37)	0.185
Smoker	8.67 (3.19-23.54)	<0.001	1.32 (0.44-3.96)	0.613
<b>Oral contraceptives</b>				
Yes	0.85 (0.63-1.14)	0.294	0.91 (0.77-1.07)	0.280
No (Ref)	1		1	
<b>Fertility drugs</b>				
Yes	1.86 (1.04-3.34)	0.036	1.51 (1.06-2.15)	0.002
No	1		1	
<b>Infertility history</b>				
Yes	1.93 (1.21-3.09)	0.005	1.35 (1.01-1.81)	0.041
Single	1.04 (0.13-8.11)	0.968	1.78 (0.71-4.46)	0.212
No (Ref)	1		1	
<b>Tubectomy</b>				
Yes	0.72 (0.50-1.02)	0.060	1.01 (0.85-1.20)	0.899
No (Ref)	1		1	
<b>Hysterectomy</b>				
Yes	7.06 (5.10-9.78)	<0.001	1.89 (1.47-2.43)	<0.001
No (Ref)	1		1	

POI: primary ovarian insufficiency, EM: early menopause, BMI: body mass index

\*In this analysis, the age at natural menopause was considered as the reference category

\*\*crude odds Ratio (95% Confidence Interval). The variables of age of menarche, BMI, gravidity, parity, and number of abortions, education, quintiles of wealth index, smoking status, fertility drug, infertility history, and tubectomy were identified

as independent variables associated with the POI in the bivariate analysis with  $P < 0.2$ .

BMI, gravidity, parity, quintiles of wealth index, smoking status, fertility drug use, and infertility history were identified as independent variables associated with early menopause in the bivariate analysis ( $P < 0.2$ )

The results of multivariable logistic regression also showed that underweight

women (BMI < 18.5) (aOR 4.12; 95% CI 1.0 to 16.9,  $P = 0.049$ ) had more than 4 times the odds of early menopause.

**Table 4.** Multivariable logistic regression analysis for factors associated with POI and EM

Variable	POI (<40)		EM (40-45)	
	aOR (95% CI)**	P-Value	aOR (95% CI)**	P-Value
<b>Age of menarche (Years)</b>				
Normal (12-14)	1			
Early <12	0.99 (0.58-1.69)	0.996	-	-
Late >14	0.67 (0.49-0.92)	0.014	-	-

Variable	POI (<40)		EM (40-45)	
	aOR (95% CI)**	P-Value	aOR (95% CI)**	P-Value
<b>BMI (kg/m<sup>2</sup>)</b>				
Normal (18.5–24.9)	1		1	
Underweight (<18.5)	3.00 (0.29-31.05)	0.357	4.12 (1.00-16.91)	0.049
Overweight (25-29.9)	0.61 (0.39-0.96)	0.033	0.97 (0.76-1.24)	0.830
Obese (≥30)	0.85 (0.56-1.28)	0.440	0.97 (0.77-1.23)	0.790
<b>Gravidity</b>				
<3 (Ref)	1		1	
≥3	1.33(0.64-2.78)	0.436	0.80 (0.51-1.25)	0.333
<b>Parity</b>				
<3 (Ref)	1		1	
≥3	0.37 (0.19-0.69)	0.002	0.92 (0.62-1.37)	0.702
<b>Number of abortions</b>				
No abortion (Ref)	1			
One	1.03 (0.71-1.48)	0.862	-	-
≥2	0.71 (0.44-1.16)	0.178	-	-
<b>Educational level</b>				
Illiterate	4.28 (0.95-19.23)	0.058	-	-
Primary school	4.25 (0.96-18.74)	0.056	-	-
Intermediate/ High school	3.02 (0.68-13.31)	0.140	-	-
University (Ref)	1			
<b>Quintiles of wealth index</b>				
1 (poorest)	1.22 (0.70-2.12)	0.475	1.21 (0.93-1.57)	0.150
2	0.91 (0.50-1.65)	0.771	1.05 (0.79-1.39)	0.714
3	1.26 (0.70-2.25)	0.430	1.02 (0.76-1.37)	0.881
4	0.97 (0.52-1.83)	0.942	0.89 (0.64-1.23)	0.491
5 (richest) (Ref)	1		1	
<b>Smoking status</b>				
Never	1		1	
Ex-smoker	1.01 (0.23-4.44)	0.987	0.56 (0.21-1.49)	0.248
Smoker	7.85 (2.60-23.67)	<0.001	1.63 (0.53-5.02)	0.390
<b>Fertility drugs</b>				
No (Ref)	1		1	
Yes	0.78 (0.29-2.05)	0.619	1.22 (0.66-2.25)	0.550
<b>Infertility history</b>				
Single	0.55 (0.06-4.53)	0.579	1.32 (0.50-3.46)	0.631
Yes	1.42 (0.64-3.16)	0.382	1.02 (0.62-1.69)	0.905
No (Ref)	1		1	
<b>Tubectomy</b>				
No (Ref)	1			
Yes	0.73 (0.50-1.07)	0.110	-	-

POI: primary ovarian insufficiency, BMI: body mass index

\*In this analysis, the age at natural menopause was considered as the reference category.

\*\*Adjusted Odds Ratio (95% Confidence Interval)

\*\*\*Significant (P < 0.05)

## Discussion

The present study determined the prevalence of POI and early menopause and identified associated factors among postmenopausal women in the AZAR cohort study in Shabestar, Iran. The results revealed that the prevalence of POI, early menopause, and age of natural

menopause is 6.1%, 27.6%, and 66.3%, respectively. Regarding the associated factors, the results indicated a statistically significant association between some socio-demographic and anthropometric characteristics and the occurrence of POI and EM.

The mean age of natural menopause in the present study was 47.4 years, consistent with

other studies conducted in the Iranian population (20, 22, 27). Generally, the mean age of natural menopause in the Middle Eastern population (46.9-47.8) is lower than in European (52.8-50.1) and American (50.5-51.4) countries (28). A reason for the difference between Iran and developed countries may be differences in lifestyle, public health, and the higher quality of health care in developed countries (29). In the present study, the prevalence of POI and early menopause was 6.1% and 27.6%, respectively. The overall prevalence of POI and early menopause varies between 1-5% and 5-10%, respectively, worldwide (30). The results of a systematic review and meta-analysis in Iran reported the prevalence of POI and early menopause at 3.7% and 12.2%, respectively (6).

Methodological differences may explain the differences in the results of the studies. In other words, the use of different approaches to estimate the prevalence of POI and early menopause in various studies can be considered the cause of this difference. Also, to evaluate the prevalence of POI, some studies determined it in the entire population, resulting in lower reported prevalence than the actual level (31, 32). The increasing prevalence of POI and early menopause in the Iranian population increases the significance of identifying the associated factors in the occurrence of these disorders.

The results of this study revealed that parity and age at menarche are associated with the occurrence of POI. In other words, the odds of POI in women with Parity $\geq$ 3 and women with late menarche (age of menarche $\geq$ 14) were 63% and 33% lower, respectively. In line with the results of our study, a study by Mishra et al. (2018) on 50,000 postmenopausal women from England, Scandinavia, Australia, and Japan showed that women who were nulliparous and had early menarche (age at menarche less than 12 years) compared to multiparous women (two or more children) and women with late menarche were at a greater risk of POI more than 5 times. In other words, early menarche alone increases the risk of POI by 80%, while this risk doubles for nulliparous women (7).

A possible explanation for the protective effect of parity against developing POI could be that ovulation stops during pregnancy. Hence,

increasing parity results in more eggs in the ovaries, which helps prevent POI in women (33). Additionally, regarding the relationship between early menopause and POI, genetic studies have indicated that some menarche-related single-nucleotide polymorphisms (SNPs) predict natural menopause (34).

Overweight was another associated factor of POI in this study (BMI 25.0-29.9). Overweight women in the present study had 39% lower odds of developing POI. Obesity is associated with reproductive function and potentially with menopause time. However, the results of studies on the relationship between obesity and menopause are very contradictory. Some studies reported the odds of POI occurrence more in underweight women (35), and some reported its occurrence more in obese women (36). However, some studies have not reported any relationship (37). Consistent with the results of the present study, the results of a prospective cohort study by Szegda et al. (2017) on 78,759 postmenopausal women showed a non-linear relationship between BMI and menopausal age. Overweight women had 21-30% lower odds of developing POI compared to normal-weight women (38). Moreover, the results of another meta-analysis revealed a moderate relationship between overweight and menopause at a later age (39). In line with the results of the present study, the study by Rizvanović et al. (2022) showed that menopausal age increases with increasing BMI (40).

In contrast with the results of the present study, the study by Rostami Dovom et al. (2021) showed that the menopause time in underweight and overweight women is approximately 0.09 and 0.03 years, respectively, shorter than in normal-weight women (21). The difference in the results of the studies may be related to their cross-sectional design. Additionally, the protective effect of overweight on POI may be mediated by estrone production in adipose tissue in overweight and obese women, as adipose tissue is a vital site for the environmental production and metabolism of estrogen in women.

In adipose tissue, circulating androstenedione produced by the ovaries and adrenal glands is converted to estrone (a weak

form of estrogen) by the aromatase enzyme. The expression and activity of aromatase in adipose tissue increase as a function of body weight and age. Estrone contributes to a woman's total estrogen levels and is positively associated with body weight. It is believed that this excess estrogen contributes to delayed menopause (41). Another possible association between BMI and menopausal age is leptin production in adipose tissue. Leptin plays a role in transmitting information about the body's energy reserves to the hypothalamus and regulating normal reproductive function. A study found that early menopause is associated with low leptin levels (42). However, more studies are needed to confirm the relationship between leptin and menopausal age.

In contrast to the two mechanisms mentioned, another mechanism proposed for the relationship between BMI and POI is that anti-mullerian hormone (AMH) is produced by granulosa cells in the ovary. It is an indicator of ovarian reserve. Some studies have indicated lower levels of AMH among obese women compared to women of normal weight and reported that obesity is associated with an increased risk of POI (43). However, this claim has not been proven well since it is still unclear what leads to a reduction in AMH levels among obese women (44). In this regard, a study by Su et al. found that AMH levels do not differ between obese and normal-weight women (45).

Smoking was another factor associated with POI. The results of the present study revealed that female smokers are almost 8 times more exposed to POI. There are several hypotheses regarding the association between smoking and early menopause. The first hypothesis is that smoking may affect ovarian aging and follicle storage by affecting gonadotropins and sex steroids, and may also have toxic effects on ovarian germ cells (46). The second hypothesis is that smoking activates aromatic hydrocarbon receptors by producing polycyclic aromatic hydrocarbons (PAHs) (47).

In this regard, many studies reported an association between smoking and POI (48, 49). In line with the results of the present study, the results of pooled analyses of 17 observational studies conducted by Zhu et al. (2018) in seven countries, including Australia, Denmark, France,

Japan, Sweden, England, and the United States, showed that current smokers are about 2 times more exposed to POI compared with never smokers. In other words, higher intensity, longer duration, higher cumulative dose, younger age at smoking initiation, and shorter time since smoking cessation were significantly associated with higher risk of POI (50). The results of the study by Vatankhah et al. (2023) in Iran also showed that women smokers had almost 2 times higher odds of POI (20).

Women's underweight (BMI < 18.5) was the only factor associated with early menopause identified in the present study. In other words, women with a BMI < 18.5 were more than 4 times more likely to experience early menopause compared to normal-weight women. Consistent with the results of our study, the results of a study by Zhu et al. (2018) revealed that underweight women (BMI < 18.5) compared to women with normal BMI (BMI = 18.5-24.9) had more than twice the odds of early menopause, and overweight and obese women were almost 1.5 times more exposed to late menopause (2). The mechanisms related to underweight and early menopause are unclear, although some hypotheses have been proposed in this regard.

The proposed hypothesis is that women with a lower BMI (BMI < 18.5) have poor fat storage, which may lead to poor-quality ovarian follicles and, thus, POI (40). Also, low body weight increases the risk of functional hypothalamic amenorrhea (FHA), which leads to anovulation, hypoestrogenism, and an increased risk of infertility. Anovulation induced by disruption of the hypothalamic-pituitary-gonadal (HPG) axis is associated with early menopause (37).

To the best of our knowledge, this study is the first study conducted to determine the prevalence of POI and early menopause and the associated factors in postmenopausal women by classifying menopausal age into three different groups in Iran. The most significant strength of the study is the use of data from a cohort study of postmenopausal women, a large sample size, and the adjustment for potential confounding factors, including demographic and reproductive data. Also, the comprehensive evaluation of women through a structured questionnaire covering many factors, such as

socio-demographic and anthropometric characteristics, and fertility history, is another strength of this study.

However, the present study suffers from some limitations. This study was cross-sectional and cannot investigate the causal relationships among variables and POI or early menopause. Also, regarding women's menopausal age, women's self-reported information about menopausal age was sufficient, which may have been subject to recall bias. Hence, to ensure menopausal age, future studies can perform hormonal tests to determine serum FSH and estradiol levels. We should treat with caution in interpreting the results obtained for smoking and underweight women due to the small sample size in these groups. The results of the study show that Iran's situation may differ from that of other countries due to differences in health system policies. Finally, more studies are needed to identify the associated factors of POI and early menopause by conducting longitudinal studies with larger sample size and following Iranian women from the early years of fertility to menopause.

## Conclusion

The results of this study showed that the prevalence of POI, early menopause, and age of natural menopause is 6.1%, 27.6%, and 66.3%, respectively. Additionally, parity, age at menarche, overweight (BMI=25.0-29.9), and smoking had a statistically significant relationship with POI. Also, a significant statistical relationship was observed between underweight (BMI<18.5) and early menopause. Since women spend more than a third of their lives in menopause, and this period is associated with unpleasant side effects for many women, health policymakers should try to enhance women's knowledge about menopause, its symptoms, and ways to improve quality of life by integrating educational programs in health centers. In addition, due to the increasing prevalence of POI, early menopause, and their associated risk factors, screening to identify the related factors by healthcare providers in health centers will be helpful in preventing and providing early interventions to increase the quality of life of women and reduce the complications caused by these disorders.

## Declarations

### Acknowledgments

The authors are grateful for financial support from the Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. The authors are also deeply indebted to all subjects who participated in this study. We appreciate the contribution by the investigators and the staff of the AZAR cohort study. We thank the close collaboration of the Shabestar health centers. In addition, we would like to thank the Persian cohort study staff for their technical support. We thank the Iranian Ministry of Health and Medical Education for contributing to the funding used in the PERSIAN Cohort through Grant no.700/534. We would like to thank the Persian Cohort Study staff for their technical support. We also express our appreciation to the Clinical Research Development Unit of Imam Reza General Hospital, Tabriz, Iran, for its cooperation, in conducting this research.

### Conflicts of interest

The authors declared no conflicts of interest.

### Ethical considerations

Informed consent was obtained from all participants before data collection. Also, a comprehensive explanation of the study's goals and process, the confidentiality of information, and the freedom to withdraw from the study at each stage was provided to all participants before the start of the study. All ethical principles in the present study were observed in accordance with the Declaration of Helsinki.

### Code of Ethics

The current study was approved by the Ethics Committee of Tabriz University of Medical Sciences (Ethical approval code: tbzmed.rec.1393.205).

### Use of Artificial Intelligence (AI)

We have not used any AI tools or technologies to prepare this manuscript.

### Funding

This study was supported by the Liver and Gastrointestinal Diseases Research Center (Grant No. 700/108 on 14 March 2016), Tabriz University of Medical Sciences. The funder had

no role in the study design, data analysis, interpretation, or writing the manuscript in this study.

### Authors' contribution

MM and EF contributed to the study design. MM and SMA contributed to the implementation and analysis plan, and MM wrote the first draft of this article. All authors critically reviewed the text, contributed in revisions, and approved the final manuscript.

### References

- Hall JE. Endocrinology of the menopause. *Endocrinology and Metabolism Clinics of North America*. 2015; 44(3): 485-496.
- Zhu D, Chung HF, Pandeya N, Dobson AJ, Kuh D, Crawford SL, et al. Body mass index and age at natural menopause: an international pooled analysis of 11 prospective studies. *European Journal of Epidemiology*. 2018; 33(8): 699-710.
- Casper R. Clinical manifestations and diagnosis of menopause-UpToDate. 2020.
- Vincent AJ, Laven JS. Early Menopause/Premature Ovarian Insufficiency. *Seminars in Reproductive Medicine*. 2020; 38(4-05): 235-236.
- Burkard T, Moser M, Rauch M, Jick SS, Meier CR. Utilization pattern of hormone therapy in UK general practice between 1996 and 2015: a descriptive study. *Menopause*. 2019; 26(7): 741-749.
- Golezar S, Ramezani Tehrani F. The global prevalence of primary ovarian insufficiency and early menopause: a meta-analysis. *Climacteric*. 2019; 22(4): 403-411.
- Mishra GD, Pandeya N, Dobson AJ, Chung HF, Anderson D, Kuh D, et al. Early menarche, nulliparity and the risk for premature and early natural menopause. *Human Reproduction*. 2017; 32(3): 679-686.
- Santoro N. Mechanisms of premature ovarian failure. *Annales d'endocrinologie*. 2003; 64(2): 87-92.
- McKinlay SM. The normal menopause transition: an overview. *Maturitas*. 1996; 23(2): 137-145.
- Panay N, Anderson RA. Premature ovarian insufficiency: an International Menopause Society White Paper. *Climacteric*. 2020; 23(5): 426-446.
- Webber L, Davies M, Anderson R, Bartlett J, Braat D, Cartwright B, et al. ESHRE Guideline: management of women with premature ovarian insufficiency. *Human Reproduction*. 2016; 31(5): 926-937.
- Xu X, Jones M, Mishra GD. Age at natural menopause and development of chronic conditions and multimorbidity: results from an Australian prospective cohort. *Human Reproduction*. 2020; 35(1): 203-211.
- Silva de Sá MF. Premature Ovarian Insufficiency and Bone Health Care: A Concern of the Gynecologist. *Revista Brasileira de Ginecologia e Obstetrícia*. 2018; 40(6): 305-308.
- Rezende GP, Dassi T. Cardiovascular Risk Factors in Premature Ovarian Insufficiency using Hormonal Therapy. *Revista Brasileira de Ginecologia e Obstetrícia*. 2023; 45(6): 312-318.
- Kulaksizoglu M, Ipekci SH, Kebapcilar L, Kebapcilar AG, Korkmaz H, Akyurek F, et al. Risk factors for diabetes mellitus in women with primary ovarian insufficiency. *Biological Trace Element Research*. 2013; 154(3): 313-320.
- Vujović S, Ivović M, Tančić Gajić M, Marina L, Jovičić SP, Pavlović N, et al. Alzheimer's Disease and Premature Ovarian Insufficiency. *Endocrines*. 2023; 4(2): 250-256.
- Blümel JE, Mezones-Holguín E, Chedraui P, Soto-Becerra P, Arteaga E, Vallejo MS. Is premature ovarian insufficiency associated with mortality? A three-decade follow-up cohort. *Maturitas*. 2022; 163: 82-87.
- Xi D, Chen B, Tao H, Xu Y, Chen G. The risk of depressive and anxiety symptoms in women with premature ovarian insufficiency: a systematic review and meta-analysis. *Archives of Women's Mental Health*. 2023; 26(1): 1-10.
- McDonald IR, Welt CK. Health-related quality of life in women with primary ovarian insufficiency: a scoping review of the literature and implications for targeted interventions. *Human Reproduction*. 2022; 37(12): 2817-2830.
- Vatankhah H, Khalili P, Vatanparast M, Ayooobi F, Esmaeili-Nadimi A, Jamali Z. Prevalence of early and late menopause and its determinants in Rafsanjan cohort study. *Scientific Reports*. 2023; 13(1): 1847.
- Rostami Dovom M, Bidhendi-Yarandi R. Prevalence of premature ovarian insufficiency and its determinants in Iranian populations: Tehran lipid and glucose study. *BMC Women's Health*. 2021; 21(1): 79.
- Zamaniyan M, Moosazadeh M, Peyvandi S, Jaefari K, Goudarzi R, MoradinAZAR M, et al. Age of Natural Menopause and Related Factors among the Tabari Cohort. *The Journal of Menopausal Medicine*. 2020; 26(1): 18-23.

23. Eghtesad S, Mohammadi Z, Shayanrad A, Faramarzi E, Joukar F, Hamzeh B, et al. The PERSIAN cohort: providing the evidence needed for healthcare reform. *The Archives of Iranian Medicine Journal*. 2017; 20(11): 691-695.
24. Poustchi H, Eghtesad S, Kamangar F, Etemadi A, Keshtkar A-A, Hekmatdoost A, et al. Prospective epidemiological research studies in Iran (the PERSIAN Cohort Study): rationale, objectives, and design. *American Journal of Epidemiology*. 2018; 187(4): 647-655.
25. Farhang S, Faramarzi E, Amini Sani N, Poustchi H, Ostadrahimi A, Alizadeh BZ, et al. Cohort Profile: The AZAR cohort, a health-oriented research model in areas of major environmental change in Central Asia. *The International Journal of Epidemiology*. 2019; 48(2): 382.
26. World Health Organization. Research on the Menopause in the 1990s: Report of a WHO Scientific Group. World Health Organization Technical Report Series. 1996: 866: 1-107.
27. Azadi T, Arghavani H, Karezani P, Sayehmiri K. Estimation of mean age of menopause in Iran: a systematic review and meta-analysis. *Journal of Ilam University of Medical Sciences*. 2018; 26(4): 85-93.
28. Palacios S, Henderson VW, Siseles N, Tan D, Villaseca P. Age of menopause and impact of climacteric symptoms by geographical region. *Climacteric : the journal of the International Menopause Society*. 2010; 13(5): 419-428.
29. Schoenaker DA, Jackson CA, Rowlands JV, Mishra GD. Socioeconomic position, lifestyle factors and age at natural menopause: a systematic review and meta-analyses of studies across six continents. *The International Journal of Epidemiology*. 2014; 43(5): 1542-1562.
30. Shuster LT, Rhodes DJ, Gostout BS, Grossardt BR, Rocca WA. Premature menopause or early menopause: long-term health consequences. *Maturitas*. 2010; 65(2): 161-166.
31. Luborsky JL, Meyer P, Sowers MF, Gold EB, Santoro N. Premature menopause in a multi-ethnic population study of the menopause transition. *Human Reproduction*. 2003; 18(1): 199-206.
32. Shamsa DS, Al Hashimi BA. Onset of natural menopause and factors affecting its development: a study from primary health care centres in Al-Risafa in Baghdad. *Iraqi New Medical Journal*. 2017; 3: 17-26.
33. Jarvis GE. Early embryo mortality in natural human reproduction: What the data say. *Research*. 2016; 5: 2765.
34. Day FR, Ruth KS, Thompson DJ, Lunetta KL, Pervjakova N, Chasman DI, et al. Large-scale genomic analyses link reproductive aging to hypothalamic signaling, breast cancer susceptibility and BRCA1-mediated DNA repair. *Nature Genetics*. 2015; 47(11): 1294-1303.
35. Hardy R, Mishra GD, Kuh D. Body mass index trajectories and age at menopause in a British birth 493 cohort. *Maturitas*. 2008; 59(4): 304-314.
36. Dratva J, Gómez Real F, Schindler C, Ackermann-Lieblich U, Gerbase MW, Probst-Hensch NM, et al. Is age at menopause increasing across Europe? Results on age at menopause and determinants from two population-based studies. *Menopause*. 2009; 16(2): 385-394.
37. Bromberger JT, Matthews KA, Kuller LH, Wing RR, Meilahn EN, Plantinga P. Prospective study of the determinants of age at menopause. *American Journal of Epidemiology*. 1997; 145(2): 124-133.
38. Szegda KL, Whitcomb BW, Purdue-Smithe AC, Boutot ME, Manson JE, Hankinson SE, et al. Adult adiposity and risk of early menopause. *Human Reproduction*. 2017; 32(12): 2522-2531.
39. Tao X, Jiang A, Yin L, Li Y, Tao F, Hu H. Body mass index and age at natural menopause: a meta-analysis. *Menopause*. 2015; 22(4): 469-474.
40. Rizvanović M, Begić Z, Kamerić L, Cerovac A, Rizvanović S, Begić M, et al. The influence of body mass index and smoking on the age of onset of menopause in women in Bosnia and Herzegovina: a cross-sectional multicentric study. *Przegląd Menopauzalny*. 2022; 21(3): 180-184.
41. Nelson LR, Bulun SE. Estrogen production and action. *Journal of the American Academy of Dermatology*. 2001; 45(3 Suppl): S116-S124.
42. Saraç F, Öztekin K, Çelebi G. Early menopause association with employment, smoking, divorced marital status and low leptin levels. *Journal of Gynaecological Endocrinology*. 2011; 27(4): 273-278.
43. Steiner AZ, Stanczyk FZ, Patel S, Edelman A. Antimüllerian hormone and obesity: insights in oral contraceptive users. *Contraception*. 2010; 81(3): 245-248.
44. Jung S, Allen N, Arslan AA, Baglietto L, Brinton LA, Eggleston BL, et al. Demographic, lifestyle, and other factors in relation to antimüllerian hormone levels in mostly late premenopausal women. *International Journal of Fertility and Sterility*. 2017; 107(4): 1012.e2-1022.e2.

45. Su HI, Sammel MD, Freeman EW, Lin H, DeBlasis T, Gracia CR. Body size affects measures of ovarian reserve in late reproductive age women. *Menopause*. 2008; 15(5): 857-861.
46. Cramer DW, Harlow BL, Xu H, Fraer C, Barbieri R. Cross-sectional and case-controlled analyses of the association between smoking and early menopause. *Maturitas*. 1995; 22(2): 79-87.
47. Blanck HM, Marcus M, Tolbert PE, Schuch C, Rubin C, Henderson AK, et al. Time to menopause in relation to PBBs, PCBs, and smoking. *Maturitas*. 2004; 49(2): 97-106.
48. Shelling AN, Ahmed Nasef N. The Role of Lifestyle and Dietary Factors in the Development of Premature Ovarian Insufficiency. *Antioxidants*. 2023; 12(8).
49. Whitcomb BW, Purdue-Smithe AC, Szegda KL, Boutot ME, Hankinson SE, Manson JE, et al. Cigarette Smoking and Risk of Early Natural Menopause. *American Journal of Epidemiology*. 2018; 187(4): 696-704.
50. Zhu D, Chung HF. Relationships between intensity, duration, cumulative dose, and timing of smoking with age at menopause: A pooled analysis of individual data from 17 observational studies. *PLOS Medicine*. 2018; 15(11): e1002704.